



ATTENTION, CONCENTRATION & INFORMATION PROCESSING

POST ACQUIRED BRAIN INJURY

Shawn Marshall MD FRCPC, Cecilia Flores-Sandoval PhD,
Penny Welch-West M.Cl.Sc. SLP, Shannon Janzen MSc, Amber
Harnett MSc RN, Connie Ferri MSc SLP, Robert Teasell MD
FRCPC

Disclaimer

This review has been prepared based on the scientific and professional information available up to July 2021. The Evidenced Based Review of Acquired Brain Injury (ERABI) information is provided for informational and educational purposes only. If you have or suspect you have a health problem, you should consult your health care provider. The ERABI contributors shall not be liable for any damages, claims, liabilities, costs, or obligations arising from the use or misuse of this material.

Copyright

With the exception of those portions of this document for which a specific prohibition or limitation against copying appears, the balance of this document may be reproduced and published in its entirety, without modification, in any form, including in electronic form, for educational or non-commercial purposes. Should any adaptation of the material be required for any reason, written permission must be obtained from ERABI. Appropriate credit or citation must appear on all copied material as follows:

Marshall S, Flores-Sandoval C, Welch-West P, Janzen S, Harnett A, Ferri C, Teasell R. (2022). Attention, Concentration, and Information Processing Post Acquired Brain Injury. In Teasell R, Marshall S, Janzen S, Cullen N, MacKenzie H, Bayley M, editors. Evidence-Based Review of Moderate to Severe Acquired Brain Injury. Version 15.0.

Funding

This work is supported by the Ontario Ministry of Health. All work produced by ERABI is editorially independent from its funding source.

Conflict of Interest

In the context of ERABI development, the term “conflict of interest” (COI) refers to situations in which an author or ERABI staff member’s financial, professional, intellectual, personal, organizational or other relationships may compromise their ability to independently conduct this evidence-based review. No limiting conflicts were identified.

Contact Information

Evidence-Based Review of Moderate to Severe Acquired Brain Injury

550 Wellington Rd South, P.O Box 5777, Stn. B., London, Ontario, Canada N6A 4V2

Website: www.ERABI.ca

Greetings from Dr. Robert Teasell,

Professor and Chair-Chief of Physical Medicine and Rehabilitation



The Collaboration of Rehabilitation Research Evidence (CORRE) team is delighted to present the Evidence-Based Review of moderate to severe Acquired Brain Injury (ERABI): *Attention, Concentration & Information Processing Post Acquired Brain Injury*. Through collaboration of researchers, clinicians, administrators, and funding agencies, ERABI provides an up-to-date review of the current evidence in brain injury rehabilitation. ERABI synthesizes the research literature into a utilizable format, laying the foundation for effective knowledge transfer to improve healthcare programs and services.

We offer our heartfelt thanks to the many stakeholders who are able to make our vision a reality. Firstly, we would like to thank the Ontario Ministry of Health, which recognizes ERABI's capacity to lead in the field of brain injury evidence-based reviews and is committed to funding it. We would also like to thank the co-chairs of ERABI, Dr. Mark Bayley (University of Toronto) and Dr. Shawn Marshall (University of Ottawa) for their invaluable expertise and stewardship of this review. Special thanks to the authors for generously providing their time, knowledge and perspectives to deliver a rigorous and robust review that will guide research, education and practice for a variety of healthcare professionals. We couldn't have done it without you! Together, we are building a culture of evidence-based practice that benefits everyone.

We invite you to share this evidence-based review with your colleagues, patient advisors that are partnering within organizations, and with the government agencies with which you work. We have much to learn from one another. Together, we must ensure that patients with brain injuries receive the best possible care every time they require rehabilitative care – making them the real winners of this great effort!

Robert Teasell, MD FRCPC

TABLE OF CONTENTS

PREFACE	6
About the Authors	6
Purpose	7
Key Concepts.....	8
Methods.....	9
Interpretation of the Evidence.....	10
Strength of the Evidence.....	11
SUMMARY OF THE EVIDENCE	13
INTRODUCTION.....	16
Non-Pharmacological Interventions	17
Drill & Practice	17
Dual-Task Training.....	19
Technological Interventions.....	20
Computer-based Interventions	20
Virtual Reality	27
Attention Training Programs.....	29
Goal Management Training	35
Mindfulness Training	37
Music Therapy.....	39
Dance Therapy	40
Animal Assisted Therapy	42
Brain Stimulation Techniques	43
Transcranial Direct Current Stimulation	43
Transcranial Magnetic Stimulation	45
Transcranial Photobiomodulation Therapy	46
Pharmacological Interventions	48

Donepezil	48
Methylphenidate	50
Bromocriptine	56
Cerebrolysin	57
Rivastigmine.....	59
Amantadine.....	61
Hyperbaric Oxygen Therapy.....	62
Dextroamphetamine	63
REFERENCES	65

PREFACE

About the Authors

ERABI is internationally recognized and led by a team of clinicians and researchers with the goal of improving patient outcomes through research evidence. Each ERABI module is developed through the collaboration of many healthcare professionals and researchers.



Dr. Shawn Marshall is a physician specializing in Physical Medicine and Rehabilitation (Physiatrist). He is the Division Head of Physical Medicine and Rehabilitation at the University of Ottawa and The Ottawa Hospital where he manages both in-patients and out-patient clinics for patients with concussion to severe traumatic brain injury. Dr. Marshall has a Master's degree in Clinical Epidemiology and is a Full Professor at the University of Ottawa in the Department of Medicine.



Cecilia Flores-Sandoval, PhD, is a clinical research assistant and the coordinator of the Evidence-Based Review of Acquired Brain Injury (ERABI). She completed a master's degree and a PhD in Health and Rehabilitation Sciences, field of Health and Aging. Her research interests include aging and rehabilitation, patient engagement and transitional care for older adults.




Penny Welch-West is a medical Speech-Language Pathologist working in Complex and Continuing Care and Acquired Brain Injury at Parkwood Institute, SJHC in London, ON. She practices in rehabilitation through palliative care in the areas of dysphagia, cognitive-communication, voice, augmentative and alternative communication. She has a special interest in supporting post secondary students in returning to school following a brain injury. She holds a Lecturer position with the University of Western Ontario and an Adjunct Clinical Professor position with McMaster University. Penny serves on many committees advocating for patient access to high quality and interdisciplinary services in



Shannon Janzen, MSc, is a research associate and the project coordinator for the Evidence-Based Review of Acquired Brain Injury (ERABI). Her research interests focus on the integration of best evidence into clinical practice to optimize patient outcomes, with an emphasis on knowledge translation initiatives.



Amber Harnett, MSc, RN, CNF scholar, completed her MSc in pathology and the accelerated BScN program at Western University. Passionate about supporting and advocating for those with acquired brain injuries, she works as a research coordinator to improve healthcare systems through research synthesis, guidelines development, knowledge translation, education, and outreach, in the CORRE lab at Parkwood Institute.



Connie Ferri is a speech-language pathologist at Parkwood Institute.



Dr. Robert Teasell is Professor of Physical Medicine and Rehabilitation, Schulich School of Medicine and Dentistry, Western University and a Clinical Researcher at Lawson Research Institute in London, Ontario. He is a clinician at Parkwood Institute, St. Joseph's Health Care London.

Purpose

The ERABI is a systematic review of the rehabilitation literature of moderate to severe acquired brain injuries (ABI). It is a regularly updated, freely accessible online resource that provides level of evidence statements regarding the strength of various rehabilitation interventions based on research studies. The

ERABI is a collaboration of researchers in London, Toronto and Ottawa. Our mission is to improve outcomes and efficiencies of the rehabilitation system through research synthesis, as well as from providing the foundational research evidence for guideline development, knowledge translation, and education initiatives to maximize the real-world applications of rehabilitation research evidence.

Key Concepts

Acquired Brain Injury

For the purposes of this evidence-based review, we used the definition of ABI employed by the [Toronto Acquired Brain Injury Network](#) (2005). ABI is defined as damage to the brain that occurs after birth and is not related to congenital disorders, developmental disabilities, or processes that progressively damage the brain. ABI is an umbrella term that encompasses traumatic and non-traumatic etiologies.

TABLE 1 | Defining Acquired Brain Injury

Included in ABI definition	Excluded from ABI definition
<p>Traumatic Causes</p> <ul style="list-style-type: none"> • Motor vehicle accidents • Falls • Assaults • Gunshot wounds • Sport Injuries <p>Non-traumatic Causes</p> <ul style="list-style-type: none"> • Tumours (benign/meningioma only) • Anoxia • Subarachnoid hemorrhage (non-focal) • Meningitis • Encephalitis/encephalopathy (viral, bacterial, drug, hepatic) • Subdural Hematoma 	<p>Vascular and Pathological Incidents</p> <ul style="list-style-type: none"> • Intracerebral hemorrhage (focal) • Cerebrovascular accident (i.e., stroke) • Vascular accidents • Malignant/metastatic tumours <p>Congenital and Developmental Problems</p> <ul style="list-style-type: none"> • Cerebral Palsy • Autism • Developmental delay • Down’s syndrome • Spina bifida with hydrocephalus <p>Progressive Processes</p> <ul style="list-style-type: none"> • Alzheimer’s disease • Pick’s disease • Dementia • Amyotrophic Lateral Sclerosis • Multiple Sclerosis • Parkinson’s disease • Huntington’s disease

Given that ‘ABI’ can have multiple definitions, studies with an ‘ABI’ population can be equally heterogeneous in terms of the sample composition. Such studies may include any combination of persons with TBI, diffuse cerebrovascular events (i.e., subarachnoid hemorrhage) or diffuse infectious disorders (i.e., encephalitis or meningitis). The vast majority of individuals with ABI have a traumatic etiology; therefore, much of the brain injury literature is specific to traumatic brain injury (TBI). The

terms ABI and TBI have been used intentionally throughout ERABI to provide more information about populations where relevant.

Moderate to Severe Brain Injury

ABI severity is usually classified according to the level of altered consciousness experienced by the patient following injury (Table 2). The use of level of consciousness as a measurement arose because the primary outcome to understand the severity of an injury is the Glasgow Coma Scale (GCS). Consciousness levels following ABI can range from transient disorientation to deep coma. Patients are classified as having a mild, moderate or severe ABI according to their level of consciousness at the time of initial assessment. Various measures of altered consciousness are used in practice to determine injury severity. Common measures include the GCS, the duration of loss of consciousness (LOC), and the duration of post-traumatic amnesia (PTA). Another factor used to distinguish moderate and severe brain injury is evidence of intracranial injury on conventional brain imaging techniques which distinguish severity of injury from a mild or concussion related brain injury.

TABLE 2 | Defining Severity of Traumatic Brain Injury, adapted from Veterans Affairs Taskforce (2008) and Campbell (2000)

Criteria	Mild	Moderate	Severe	Very Severe
Initial GCS	13-15	9-12	3-8	Not defined
Duration LOC	< 15minutes*	<6 hours	6-48 hours	>48 hours
Duration PTA	< 1hour*	1-24 hours	1-7 days	>7 days
	*This is the upper limit for mild traumatic brain injury; the lower limit is any alteration in mental status (dazed, confused, etc.).			

Methods

An extensive literature search using multiple databases (CINAHL, PubMed/MEDLINE, Scopus, EMBASE, and PsycINFO) was conducted for articles published in the English language between 1980–July 2021 that evaluate the effectiveness of any intervention/treatment related to ABI. The references from key review articles, meta-analyses, and systematic reviews were reviewed to ensure no articles had been overlooked. For certain modules that lacked research evidence the gray literature, as well as additional databases, were searched in order to ensure the topic was covered as comprehensively as possible.

Specific subject headings related to ABI were used as the search terms for each database. The search was broadened by using each specific database’s subject headings, this allowed for all other terms in the database’s subject heading hierarchy related to ABI to also be included. The consistent search terms used were “head injur*”, “brain injur*”, and “traumatic brain injur*”. Additional keywords were used specific to each module. A medical staff librarian was consulted to ensure the searches were as comprehensive as possible.

Every effort was made to identify all relevant articles that evaluated rehabilitation interventions/treatments, with no restrictions as to the stage of recovery or the outcome assessed. For each module, the individual database searches were pooled, and all duplicate references were removed. Each article title/abstract was then reviewed; titles that appeared to involve ABI and a treatment/intervention were selected. The remaining articles were reviewed in full.

Studies meeting the following criteria were included: (1) published in the English language, (2) at least 50% of the study population included participants with ABI (as defined in Table 1) or the study independently reported on a subset of participants with ABI, (3) at least three participants, (4) $\geq 50\%$ participants had a moderate to severe brain injury (as defined in Table 2), and (5) involved the evaluation of a treatment/intervention with a measurable outcome. Both prospective and retrospective studies were considered. Articles that did not meet our definition of ABI were excluded.

Interpretation of the Evidence

The levels of evidence (Table 3) used to summarize the findings are based on the levels of evidence developed by Sackett et al. (2000). The levels proposed by Sackett et al. (2000) have been modified; specifically, the original ten categories have been reduced to five levels. Level 1 evidence pertains to high quality randomized controlled trials (RCTs) (PEDro ≥ 6) and has been divided into two subcategories, level 1a and level 1b, based on whether there was one, or more than one, RCT supporting the evidence statement.

The evidence statements made in evidence-based reviews are based on the treatment of groups rather than individuals. There are times when the evidence will not apply to a specific case; however, the majority of patients should be managed according to the evidence. Ultimately, the healthcare professional providing care should determine whether an intervention is appropriate and the intensity with which it should be provided, based on their individual patient's needs. Furthermore, readers are asked to interpret the findings of studies with caution as evidence can be misinterpreted. The most common scenario occurs when results of a trial are generalized to a wider group than the evidence allows. Evidence is a tool, and as such, the interpretation and implementation of it must always be done with the known limitations in mind.

TABLE 3 | Levels of Evidence

Level	Research Design	Description
1A	Randomized Controlled Trial (RCT)	More than one RCT with PEDro score ≥ 6 . Includes within subject comparisons, with randomized conditions and crossover designs
1B	RCT	One RCT with PEDro ≥ 6
2	RCT	One RCT with PEDro < 6
	Prospective Controlled Trial (PCT)	Prospective controlled trial (not randomized)
	Cohort	Prospective longitudinal study using at least two similar groups with one exposed to a particular condition
3	Case Control	A retrospective study comparing conditions including historical controls
4	Pre-Post Trial	A prospective trial with a baseline measure, intervention, and a post-test using a single group of subjects
	Post-test	A prospective intervention study using a post intervention measure only (no pre-test or baseline measurement) with one or more groups
	Case Series	A retrospective study usually collecting variables from a chart review
5	Observational study	Using cross sectional analysis to interpret relations
	Clinical Consensus	Expert opinion without explicit critical appraisal, or based on physiology, biomechanics or "first principles"
	Case Reports	Pre-post or case series involving one subject

Strength of the Evidence

The methodological quality of each RCT was assessed using the Physiotherapy Evidence Database (PEDro) rating scale developed by the Centre for Evidence-Based Physiotherapy in Australia (Moseley et al., 2002). The PEDro is an 11-item scale; a point is awarded for ten satisfied criterion yielding a score out of ten. The first criterion relates to external validity, with the remaining ten items relating to the internal validity of the clinical trial. The first criterion, eligibility criteria, is not included in the final score. A higher score is representative of a study with higher methodological quality.

**ATTENTION,
CONCENTRATION &
INFORMATION PROCESSING**

POST ACQUIRED BRAIN INJURY

SUMMARY OF THE EVIDENCE

Intervention	Key Points Level of Evidence
Non-Pharmacological Interventions	
Drill & Practice	<p>Drill and practice training may not be effective for the remediation of attention following an ABI.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence that drill, and practice training may not be effective for the remediation of attention compared to spontaneous recovery, regardless of the level of structure in the program for those with an ABI.</i>
Dual-Task Training	<p>Dual-task training has been shown to improve measures of attention task performance when compared to non-specific/generalized training.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence that dual task training may be effective in improving attention task performance in ABI populations compared to more generalized training.</i>
Technological Interventions	<p>Computer-based may not be more effective than no intervention in improving measures of attention and concentration post ABI. However, a computer-assisted attention retraining program may improve attention in individuals with ABI, when compared to memory training or recreational computing.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence that compensatory strategies may result in better performance in attention tasks during computer-based training for individuals with ABI.</i> - <i>There is level 2 evidence that THINKable, a computer-based multi-media program is effective for improving attention in individuals with ABI.</i> - <i>There is level 1b evidence that a computer-assisted attention retraining program may be more beneficial for individuals with ABI than memory training or recreational computing, when compared.</i> - <i>There is level 1b evidence that video game play may not improve sustained attention in individuals with ABI.</i> - <i>There is level 2 evidence that rehabilitation interventions using computer programs and software have limited effects on attention in individuals with ABI.</i> <p>Repetitive tasks in virtual reality environments are effective in improving attention and concentration in individuals with ABI.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence that exercising in a virtual environment may improve concentration and reaction time but may not be effective in improving attention in individuals with ABI.</i> - <i>There is level 4 evidence that attention performance can be improved in individuals with ABI through repetition of task in virtual reality environments.</i>

<p>Attention Training Programs</p>	<p>Attention process training may improve attentional functioning in individuals with ABI. Time pressure management might improve attention performance and information processing speed.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence that adaptive training is no more effective than non-adaptive training in remediating attention in ABI populations.</i> - <i>There is level 1b evidence that the Short-Term Executive Plus (STEP) cognitive rehabilitation program is not effective in treating attentional disorders compared to waitlist controls in ABI populations</i> - <i>There is level 2 evidence that providing an external inhibitor may improve attention switching in ABI populations.</i> - <i>There is level 1a evidence that attention process training (APT) may improve attentional functioning in individuals with ABI.</i> - <i>There is level 2 evidence that time pressure management training may improve attention performance and information processing speed.</i> - <i>There is level 4 evidence that cognitive pragmatic training may not be effective for improving attention in individuals with ABI.</i> - <i>There is level 4 evidence that a working memory training program may remediate attention in individuals post ABI.</i> <p>Goal management training is effective in assisting those who sustain an ABI in learning to manage life goals through improved attention.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence to suggest goal management training, when compared to education, may be effective at improving attention in individuals post ABI.</i> - <i>There is level 2 evidence that goal management training is more effective in remediating task completion times than motor skill training; however, it is not more effective in treating attention deficits, in individuals post ABI.</i> <p>Mindfulness training may offer benefits for individuals with TBI in terms of attention; however, the length of exposure needs to be carefully considered.</p> <ul style="list-style-type: none"> - <i>There is level 2 evidence that mindfulness training compared to no intervention may improve an individual's ability to correctly reject inappropriate stimuli post ABI.</i> - <i>There is level 2 evidence that an attentional control training intervention involving mindfulness meditation may not be effective to treat attentional difficulties in individuals with TBI.</i>
<p>Music Therapy</p>	<p>Music therapy might improve cognitive function, such as attention, in individuals with severe to moderate TBI but more studies are needed.</p> <ul style="list-style-type: none"> - <i>There is level 1a evidence that neuromuscular music therapy may lead to changes in functional neuroplasticity, enhancing cognitive function, such as attention, in a population with moderate to severe TBI; however, further studies are needed.</i>
<p>Dance Therapy</p>	<p>Dance therapy may not improve attention in individuals with severe TBI.</p> <ul style="list-style-type: none"> - <i>There is level 1a evidence that a dance program may not improve attention in individuals with moderate to severe TBI.</i>
<p>Animal Assisted Therapy</p>	<p>Animal assisted therapy may improve attention span and concentration in individuals with ABI.</p>

	<ul style="list-style-type: none"> - <i>There is level 1b evidence that animal therapy may enhance attention span and concentration in individuals with an ABI.</i>
<p>Brain Stimulation Techniques</p>	<p>Transcranial direct current stimulation (tDCS) may be effective in remediating attentional deficits when combined with computer-assisted training in ABI populations.</p> <ul style="list-style-type: none"> - <i>There is level 1b evidence that participants who received active tDCS presented slower and more variable reaction time than participants in the sham group. The effects of tDCS are subtle and unreliable in terms of enhancing processing speed and working memory accuracy in individuals with ABI</i> - <i>There is level 2 evidence that tDCS when combined with a computer assisted training program (compared to sham stimulation) may improve divided attention in individuals post ABI.</i> - <i>There is level 4 evidence that tDCS may be feasible and safe for individuals with TBI.</i> <p>Repeated magnetic transcranial stimulation (rTMS) needs to be further explored; current studies have reported conflicting results as to its effectiveness in remediating attentional deficits following ABI.</p> <ul style="list-style-type: none"> - <i>There is conflicting level 1b evidence as to whether rTMS compared to sham stimulation may improve attention following an ABI.</i> <p>Transcranial photo biomodulation therapy may not be effective in improving attention in individuals with ABI.</p> <ul style="list-style-type: none"> - <i>There is level 4 evidence that transcranial photo biomodulation therapy may not improve attention following an ABI.</i>
<p>Pharmacological Interventions</p>	
<p>Donepezil</p>	<p>It is unclear as to whether donepezil may improve attention in individuals with moderate to severe ABI.</p> <ul style="list-style-type: none"> - <i>There is conflicting level 1b (positive) and level 2 (negative) evidence that donepezil may improve attention compared to placebo post ABI.</i>
<p>Methylphenidate</p>	<p>Studies examining methylphenidate for attention and information processing show conflicting results. Response to methylphenidate may depend on the presence of the Met genotype and/or dopamine transporter levels, as well as on the age of the individual.</p> <ul style="list-style-type: none"> - <i>There is conflicting level 1a evidence regarding the effectiveness of methylphenidate following brain injury for the improvement of attention in individuals post TBI.</i> - <i>There is level 1b evidence that individuals carrying the Met allele may be more responsive to methylphenidate than those without the Met allele when it comes to the TBI population.</i>
<p>Bromocriptine</p>	<p>The effectiveness of Bromocriptine on attention for individuals with ABI is unknown at this time, further studies are needed.</p> <ul style="list-style-type: none"> - <i>There is conflicting evidence (Level 1b and Level 2) evidence as to whether bromocriptine improves performance on attention tasks compared to placebo in individuals post TBI.</i>

Cerebrolysin	<p>Cerebrolysin may be beneficial for improving attentional functioning following brain injury; however, controlled trials are needed to further evaluate its efficacy.</p> <ul style="list-style-type: none"> - <i>There is level 1b evidence that cerebrolysin is an effective therapy that may improve attentional function for individuals living with moderate to severe TBI.</i>
Rivastigmine	<p>Rivastigmine may not be effective in improving concentration or processing speed post TBI.</p> <ul style="list-style-type: none"> - <i>There is level 1a evidence that Rivastigmine compared to placebo is not effective for improving concentration or processing speed in post ABI individuals but may increase vigilance.</i>
Amantadine	<p>Amantadine may not be effective in treating attention deficits following an ABI.</p> <ul style="list-style-type: none"> - <i>There is level 1b evidence that amantadine is not effective for improving attention compared to placebo following an ABI.</i>
Hyperbaric Oxygen Therapy	<p>Hyperbaric oxygen therapy may improve attention and processing speed following an ABI; however, more research is required.</p> <ul style="list-style-type: none"> - <i>There is level 4 evidence that hyperbaric oxygen therapy may improve both attention and processing speed following an ABI.</i>
Dextroamphetamine	<p>Dextroamphetamine may not be an effective treatment for attentional deficits following an ABI and may increase agitation.</p> <ul style="list-style-type: none"> - <i>There is level 1b evidence that dextroamphetamine does not improve attention following an ABI.</i>

INTRODUCTION

Attention refers to the individual’s ability to select information from a multitude of sensory input available, while filtering the unattended input (Hanh et al., 2008). Attention is usually measured using externally directed tests, such as instructing participants to focus their attention on a sequence of stimuli or attenuating to a particular stimulus. Attention has four components: working memory, competitive selection, top-down sensitive control, and automatic filtering for salient stimuli (Knudsen, 2007). The mechanisms of attention select the information that gains access to working memory, where it is analyzed in detail, allowing the individual to make decisions and plans of actions about that information (Knudsen, 2007). The attention clinical model proposed by Sohlberg and Mateer (2001) was based upon the attentional deficits experienced by individuals with TBI and includes five components of attention.

TABLE 4 | Clinical model of attention, adapted from Sohlberg and Mateer (2001)

Component	Description
Focused Attention	Refers to a response to discrete visual, auditory, or tactile stimuli (e.g., pain, temperature)
Sustained Attention	Is the ability to maintain a consistent behavioural response during continuous and repetitive activity, involves two subcomponents: Vigilance and working memory
Selective Attention	Refers to the ability to ignore irrelevant or distracting stimuli, involves the notion of ‘freedom from distractibility’ (e.g., prepare a meal with children playing in the background)
Alternating Attention	Set shifting and mental flexibility. It is the ability to shift focus of attention and move between tasks that have different cognitive requirements, and controlling which information will be processed (e.g., taking notes during a lecture)
Divided Attention	Refers to the ability to respond to multiple, simultaneous tasks. Two or more behavioural responses or stimuli may be required or need to be monitored (e.g., driving a car while listening to the radio)

Attention deficits have been reported as the most common difficulty and the most debilitating issue for individuals with TBI; often manifested during activities of daily living and related to slowed processing speed, distractibility, inability to sustain attention, and difficulties shifting attention (Roitsch et al., 2019). In general, individuals with TBI demonstrate significant deficits compared to control populations. The former have challenges in speed of information processing tasks (Dymowski et al., 2015), increased errors on dual task experiments for sustained attention (Dockree et al., 2006; Hasegawa & Hoshiyama, 2009), slower reaction times (Azouvi et al., 2004; Stuss et al., 1989), and increased mental effort on attentional tasks (Azouvi et al., 2004). However, in a case series study, Foley et al. (2010) found that level of injury severity, as measured by the GCS or PTA, did not play a role in who performed poorly on the dual task assignments. The authors found that only 27% of study participants with TBI performed below the cut-off for normal performance (Foley et al., 2010). In a recent systematic review and meta-analysis, Walz et al. (2021) found that visual attention is significantly impacted in individuals with TBI, particularly the ability to allocate attention when using top-down attentional control with a directional cue, and difficulties disengaging from an incorrect spatial cue.

Non-Pharmacological Interventions

Drill & Practice

The following studies examined the influence of “drill & practice” exercises (either computerized and/or paper-and-pencil) on attentional functioning. Drill and practice training targets attention skills through repetitive training of specific tasks involving attention.

TABLE 5 | The Effect of Drill and Practice on Attention Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
Novack et al. (1996)	Population: Severe TBI; <i>Focused Stimulation Group (n=22)</i> : Mean Age=28.7yr; Mean Time	1. Analysis of primary outcome measures revealed no significant differences between the

<p>USA RCT PEDro=5 N=44</p>	<p>Post Injury=5.9wk. <i>Unstructured Stimulation Group (n=22)</i>: Mean Age=26.4yr; Mean Time Post Injury=6.4wk. Intervention: Participants were randomly placed into a focused or unstructured stimulation group. Participants in the focused group received hierarchical attentional learning training (30min, 5x/wk). Skills were not taught in a hierarchical or sequential fashion in the unstructured group. Outcome Measures: Digit Span and Mental Control subtests of Wechsler Memory Scale-Revised (WMS-R), computer-based simple and choice reaction time tests, Logical Memory I & II, Sentence Repetition, Judgment of Line Orientation, Trail Making A & B, Arithmetic subtest Wide Range Achievement Test-Revised, Visual imperceptions.</p>	<p>focused and unstructured stimulation groups, both at baseline and discharge (WMS-R digit span and mental control, computer-based simple and choice reaction times test).</p> <ol style="list-style-type: none"> Both groups show significant improvement on attentional skills as measure by the Digit Span and Mental Control subtests of the WMS-R, and by measures of simple and choice reaction time. There were no significant differences between the groups with respect to any secondary outcome measures studied.
<p>Lindelov et al. (2016) Denmark PCT N_{initial}=78, N_{Final}=35</p>	<p>Population: <i>ABI Group (n=17)</i>: Mean Age=56.1yr; Gender: Male=13, Female=4; Mean Time Post Injury=57d. <i>Healthy Group (n=18)</i>: Mean Age=56.1yr; Gender: Male=8, Female=10. Intervention: ABI and healthy participants were randomized and analyzed separately. Experimental group participants received 20 sessions of N-back training (N-back), where participants press a key when presented stimulus is identical to the stimulus N back in the sequence. Control group participants received 20 sessions of visual search training (VS), where participants press a key if a target symbol is present in an NxN array of symbols. Outcome Measures: Raven’s Advanced Progressive Matrices (RAPM), Wechsler Adult Intelligence Scale-IV (WAIS-IV), Working Memory Index (WMI Index, digit span, arithmetic, letter-number sequencing), Operation Span Test (OSPAN), WAIS-IV Processing Speed Index (PSI index, search, coding), Stroop Test.</p>	<ol style="list-style-type: none"> Both ABI and healthy groups showed significant improvement post-intervention on the assigned training tasks. The standardized mean difference was 0.45 for ABI N-back, 6.11 for healthy N-back, 1.06 for ABI VS, and 3.34 for Healthy VS. The healthy group showed greater improvement than the ABI group. No significant differences in improvements between N-back and VS treatments (time x treatment interaction) were found in either group for WMI-digit span, WMI-arithmetic, WMI-letter-number sequencing, WMI index, PSI-search, PSI-coding, PSI index, RAPM, OSPAN, or Stroop. No significant differences in improvement between healthy and ABI groups (group x time x test interaction) were found for WMI-digit span, WMI-arithmetic, WMI-letter-number sequencing, WMI index, PSI-search, PSI-coding, PSI index, RAPM, OSPAN, or Stroop.

Discussion

The two studies reported no significant differences between groups for attentional, functional, and/or cognitive skills assessed (Lindelov et al., 2016; Novack et al., 1996). Novack et al. (1996) compared focused hierarchical attentional learning with an unstructured non-sequential, non-hierarchical intervention, while Lindelov et al. (2016) compared N-back training with visual search training. Novack et al. (1996) found that there were no significant differences between groups at either time points; however, both groups significantly improved over time. Although Lindelov et al. (2016) found no significant treatment effects over time, and no spontaneous recovery effects were observed. Overall,

there is limited evidence in support of “drill & practice” as an effective rehabilitation intervention for attention in individuals with ABI.

Conclusions

There is level 2 evidence that drill and practice training may not be effective for the remediation of attention compared to spontaneous recovery, regardless of the level of structure in the program for those with an ABI (Lindelov et al., 2016; Novack et al., 1996).



KEY POINTS

- Drill and practice training may not be effective for the remediation of attention following an ABI.

Dual-Task Training

For individuals with TBI, performing an activity in isolation may be possible; however, concurrent performance of complex cognitive tasks may be more challenging (Zavoronkova et al., 2016). Dual-task training involves dividing attention between two stimuli to complete two tasks concurrently and successfully, such as walking while speaking. The following studies examined the effect of “dual-task” training on speed of processing.

TABLE 6 | The Effect of Dual-Task Training on Speed of Processing Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
<p>Couillet et al. (2010) France RCT PEDro=5 N=12</p>	<p>Population: Severe TBI; Gender: Male=9, Female=3. <i>Group 1 (n=5):</i> Mean Age=23.8yr; Mean GCS=4.8; Mean Time Post Injury=6.3mo. <i>Group 2 (n=7):</i> Mean Age=26.7yr; Mean GCS=4.8; Mean Time Post Injury=16.1mo. Intervention: Randomized AB versus BA design, where “A” represents the control phase and “B” represents the treatment (dual-task training) phase. In the dual-task phase, participants were trained to conduct two concurrent tasks simultaneously. Group 1 started with the control phase (AB) and Group 2 (BA) with the treatment phase. Each phase lasted 6wk (4, 1 hr sessions/wk). Outcome Measures: Test Battery for Attentional Performance (TAP: divided attention and flexibility subtests), Go-no go, Digit Span, Trail Making Test, Stroop Test,</p>	<ol style="list-style-type: none"> 1. Following training, there was a significant improvement in the 2 tasks that targeted divided attention (TAP-divided attention, Go-no go and Digit Span: p<0.0001 for both). Performance was improved in both speed of response and accuracy. 2. The two groups differed significantly at 6wk, with those in the BA design doing better on TAP reaction times (p<0.01), the digit span dual-task (p<0.001), and the Rating Scale of Attentional Behaviour (p<0.01). 3. There were significant differences between groups at 6wk on the Stroop test (p<0.001) and the flexibility subtest of the TAP (p<0.001), but not the Trail Making Test or the Brown-Peterson task. 4. Experimental training had no significant effects on non-target measures.

<p>Stablum et al. (2000) Italy Case-Control N=38</p>	<p>Brown-Peterson Paradigm, Rating Scale of Attentional Behaviour.</p> <p>Population: Chronic Head Injury (CHI)=10, Gender: Male=9, Female=1; mean age:25.6yr, time since injury: 27.8 months. Controls=10. Intervention: Both participants with CHI and controls were assessed, received treatment, and were retested at 3mo follow-up. Treatment consisted of five sessions, once a week, with the dual-task condition. Retest included a measure of speed of controlled information processing. Outcome Measures: Wisconsin Card Sorting Task (WCST), Paced Auditory Serial Addition Task (PASAT), Dual-Task Paradigm.</p>	<ol style="list-style-type: none"> 1. Significant difference between participants and controls on number of preservative errors ($p<0.017$) and categories ($p<0.020$) achieved in WCST, and PASAT mean time ($p=0.031$). 2. Reaction time was slower for CHI participants than controls in dual-task ($p<0.005$); dual task cost significantly greater for CHI patients than controls ($p<0.028$). 3. At retest and at 3-months follow-up reaction time was slower for CHI participants than controls ($p<0.0001$); but participants demonstrated a greater reduction in dual-task cost after treatment (54 vs 22 ms).
--	--	---

Discussion

A single RCT demonstrated that attention and information processing outcomes could be improved within the dual-task paradigm (Couillet et al., 2010). The authors found that dual-task training significantly improved attentional behaviour and reaction time compared to a generalized cognitive program (Couillet et al., 2010). In a case control study, Stablum et al. (2000) found that individuals with a closed head injury (CHI) performed poorly on dual-task measures initially; however, with additional training, their completion time of dual-task measures significantly increased compared to the control group.

Conclusions

There is level 2 evidence that dual task training may be effective in improving attention task performance in ABI populations compared to more generalized training (Couillet et al., 2010; Stablum et al., 2000).



KEY POINTS

- Dual-task training has been shown to improve measures of attention task performance when compared to non-specific/generalized training.

Technological Interventions

Computer-based Interventions

A surge in technology has allowed for the development of more computer-based intervention solutions designed to improve attention, concentration, and information processing. Current treatment modalities include computer cognitive training programs and virtual reality sessions. Computerized

cognitive training is a low cost, user-friendly and accessible intervention that may improve cognition in individuals with ABI; additionally, computer interventions can be relatively easy implemented in a variety of settings (Sigmundsdottir et al., 2016).

TABLE 7 | The Effect of Computer-Based Interventions on Attention and Reaction Time Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
<p>Dirette et al. (1999) USA RCT PEDro=4 N=30</p>	<p>Population: TBI: Mean age=38yr; Gender: male=22, female=8; Time since injury range=2-12 months. Intervention: Randomly assigned to remedial (without instruction, n=15) and compensatory strategy (verbalization, chunking and pacing) intervention (n=15) groups receiving a 45-minute session once a week for 4 weeks. Outcome Measures: Pre and Post-test on the Paced Auditory Serial Addition Task (PASAT).</p>	<ol style="list-style-type: none"> 1. Pre/post and weekly computer-based tasks significantly improved in both groups ($p<0.01$). 2. No significant improvement due to intervention ($p>0.05$).
<p>Ruff et al. (1994) USA RCT PEDro=3 N=15</p>	<p>Population: Severe head injury; Mean Age=26.9yr; Time Post Injury\geq6mo. Intervention: Participants were randomized to one of two treatment conditions: attention training followed by memory training (Group A; n=7) or vice versa (Group B; n=8). Training was provided from THINKable, a computer-based multi-media program. Training was terminated after either 20 hr (2hr/d) were completed, or 90% scores were achieved on the most advanced program. Participants were assessed before, during and after training. Outcome Measures: 2 + 7 Selective Attention Test, WAIS-R Digit Symbol, Continuous Performance Test (CPT); Rey Auditory Verbal Learning Test, Corsi Block Learning Test.</p>	<ol style="list-style-type: none"> 1. Computer-based attention training resulted in significant improvements for attention ($p=0.003$). 2. Significant improvement in Memory II ($p=0.021$) but not Memory I or III. Gains were significant for Rey Verbal ($p=0.004$) and Corsi Block Learning ($p=0.03$) total correct as well. 3. Significant improvements in digital symbol scores ($p<0.001$) were noted as well, but no significant changes were found with CPT or 2+7 test scores.
<p>Gray et al. (1992) UK RCT PEDro=5 N=31</p>	<p>Population: Close Head Injury=17; Others=14. <i>Experimental Group (n=17):</i> Mean Age=26.18yr; Gender: Male=12, Female=5; Mean Time Post Injury=79wk. <i>Control Group (n=14):</i> Mean Age=34.14yr; Gender: Male=10, Female=4; Mean Time Post Injury=84wk. Intervention: Participants in the experimental group received micro-computerized attentional training (1-1.5hr sessions for 3-9wk). The training covered reaction time training, rapid number comparison, digit symbol transfer, and divided attention tasks. The control group received recreational computing for a similar time period. Outcome Measures: Digit Span, Backward Digit Span, Paced Auditory Serial Addition task (PASAT), Information Processing Rate (IRP),</p>	<ol style="list-style-type: none"> 1. At post-test assessment, the experimental group showed significant improvement on the WAIS-R picture completing ($p=0.031$) and the PASAT information processing rate ($p=0.023$). 2. At the 6-mo follow-up, differences between the groups indicated significant improvement on the Backward Digit Span ($p=0.007$), the WAIS-R Arithmetic ($p=0.014$), information processing rate and the PASAT ($p=0.011$), longest string ($p=0.009$), IPR ($p=0.019$). 3. For the experimental group, improvements from the intervention were found for IPR ($p=0.004$). 4. In general, course improvement was seen in the experimental group during the intervention phase and was continued into follow-up.

	<p>Longest string, Wisconsin Card Sorting Test, Wechsler Adult Intelligence Scale-Revised (WAIS-R) Arithmetic.</p>	
<p>Niemann et al. (1990) United States RCT PEDro=7 N=29</p>	<p>Population: <i>Attention Group (N=13):</i> Mean age=28.9yr; Mean time post-injury=41mo. <i>Memory Group (N=13):</i> Mean age=34.3yr; Mean time post-injury=37.1mo. Intervention: Individuals were randomly assigned to either a computer-assisted attention retraining program or a memory training program. Both programs lasted 9 weeks and had two 2-hour sessions each week. Outcomes: Attention Test d2, Paced Auditory Serial-Addition Task (PASAT), Divided Attention test (DAT), Trail Making Test-B (TMT-B), Rey Auditory Verbal Learning Test (RAVLT), Block Span Learning Test (BSLT), Ruff 2 & 7 Test, Logical Memory Subtest (Wechsler Memory Scale) (WMS-LM), Ruff-Light Trail Learning Test (RLTL).</p>	<ol style="list-style-type: none"> 1. There were no significant within-group differences on the Test d2, PASAT, DAT, RAVLTBSLT, Ruff 2 & 7 Tests, WMS-LM, or the RLTL. 2. Significant within group differences were seen on the TMT-B for both the attentional (p<0.01), and memory (p<0.01) groups. 3. The attention group improved significantly more on the TMT-B compared to the memory group (p=0.05). 4. The attention group improved significantly more than the memory group on the Attention Test d2 (p=0.02). 5. No other significant differences were found.
<p>Malec et al. (1984) United States RCT Crossover PEDro=8 N=10</p>	<p>Population: Mean age=30yr; Gender: Male=8, Female=2; Mean time post injury=80dys. Intervention: Individuals played two types of first-person shooter video games, one with no interfering targets and one with them present. Individuals were randomly assigned to treatment order. Video game conditions were 1 week-long and included twice daily sessions of video game play. Outcome Measures: Stroop Test, Letter Cancellation task, Symbol Cancellation task, reaction time (RT).</p>	<ol style="list-style-type: none"> 1. No significant differences were found between conditions at any time points.
<p>Lesniak et al. (2020) Poland PCT N= 15</p>	<p>Population: TBI; Mean Age= 26.2±7.6yr; Gender: Male=11, Female=4; Mean time post injury= 11.6±6.6mo; Severity: Severe=10, Moderate=5. Intervention: Cognitive rehabilitation therapy program focused on memory and attention. The individual therapy program was cognitive training conducted with computer software (RehaCom) and supervised by a psychologist. Group sessions were run by a neuropsychologist and focused on internal memory strategies and external aids. Participants had 15 group session (45 min, 5d/wk) and 15 individual therapy sessions (45min, 5 d/wk). Assessments were conducted at baseline (3wk prior to start), pre-treatment, posttreatment and at 4mo follow-up. Outcome Measures: Cambridge Neuropsychological Test Automated Battery (CANTAB), Pattern Recognition Memory Test, Rey’s Auditory Verbal Learning Test (RAVLT), Spatial Span Test (SSP), Paced Auditory Serial</p>	<ol style="list-style-type: none"> 1. From baseline to preintervention only the PASAT changed significantly (p=0.047). 2. From baseline to post-intervention there were no significant changes in short-term verbal memory (RAVLT; p=0.242), short-term visual memory (PRM; p=0.172) or visuospatial working memory (SSP; p=0.24). 3. From baseline to post-intervention RVP attention test (p=0.002) and PASAT (p=0.005) showed significant improvement. 4. Pre to Post intervention significant improvements were found for PRM (p=0.022), RVP (p=0.002) and PASAT (p=0.012). 5. Post-intervention, participants reported less everyday cognitive problem than at baseline (EBIQ). No significant differences were found between post-intervention and follow-up.

	Addition Test (PASAT), Rapid Visual Information Processing, European Brain Injury Questionnaire (EBIQ).	
<p>O'Neil-Pirozzi and Hsu (2016) PCT N_{Initial}=14 N_{Final}=12</p>	<p>Population: TBI=4, CVA=2, Brain tumor=1; Severity: moderate/severe. <i>Experimental Group (n=7):</i> Mean Age=51.3yr; Gender: Male=5, Female=2; Mean Time Post Injury=20.9yr; Etiology: TBI=5, CVA=2. <i>Control Group (n=7):</i> Mean Age=46.9yr; Gender: Male=7; Mean Time Post Injury=25.0yr.</p> <p>Intervention: Experimental group participants received BrainHQ, a commercially available online computerized cognitive exercise program (Attention, Brain Speed, Memory, People Skills, Intelligence, and Navigation) for 5 mo, 5d/wk. Control group participants did not have a private computer and received no intervention.</p> <p>Outcome Measures: Number/percentage of sessions completed, Number/percentage of sessions initiated by participants, Number/percentage of sessions completed independently by participants, Mean amount of external cues provided for session completion, Wisconsin Card Sorting Test (WCST), Hopkins Verbal Learning Test-Revised (HVLTR immediate, delayed), Controlled Oral Word Association Test-FAS (COWAT), Trail Making Test (TMT A and B accuracy and speed), Satisfaction with Life Scale (SWLS), Semi-structured interview questions.</p>	<ol style="list-style-type: none"> Of the five experimental group participants that completed the study, they completed an average 87% of sessions, initiated an average 25% of sessions, and independently completed an average 7% of sessions. Two participants needed minimum external cues, two participants needed moderate external cues, and one participant needed maximum external cues. Comparing 3mo prior to intervention with 1wk prior to intervention, there were no significant differences within either group for WCST, HVLTR, COWAT, TMT A or B, or SWLS. There were no significant differences between groups at 1wk prior to intervention (baseline) for WCST, HVLTR, COWAT, TMT A or B, or SWLS. Compared to baseline, experimental group showed significant improvement post-intervention for HVLTR-immediate (p=0.0255) and SWLS (p=0.0075). There were no significant improvements for WCST, HVLTR-delayed, or TMT A or B. Compared to baseline, control group did not show significant differences post-intervention for WCST, HVLTR, TMT A or B, or SWL. Compared to control group, experimental group showed significantly higher post-intervention improvements on HVLTR-immediate (p=0.0068) and COWAT (p=0.0310). No significant differences between groups were found for changes in WCST, HVLTR-delayed, TMT A or B, or SWL. Of the experimental group participants who completed the study, 60% reported improved everyday thinking abilities, 60% reported improved memory, and 20% reported improved attention, organization, and/or problem-solving skills, but 60% reported they would not continue with exercise program post-study completion.
<p>Li et al. (2015) USA Pre-Post N_{Initial}=13, N_{Final}=12</p>	<p>Population: Stroke=5, TBI=5, Brain tumor=2; Mean Age=61yr; Gender: Male=10, Female=2.</p> <p>Intervention: Participants received the computer-based cognitive retraining program, Parrot Software. The following eight modules were each completed in separate 1h sessions: Visual Instructions, Attention Perception and Discrimination, Concentration, and Visual Attention Training, Remembering Written Directions, Remembering Visual Patterns,</p>	<ol style="list-style-type: none"> Compared to baseline, there was a significant mean increase in overall MoCA of 3.25 (p=0.03) post-intervention. However, the attention and memory subscales did not show significant differences. There were no significant differences before and after intervention for the medication-box sorting task. Participants with previous computer-based cognitive retraining experience had significantly

ATTENTION, CONCENTRATION & INFORMATION PROCESSING POST ACQUIRED BRAIN INJURY

	<p>Remembering Written Letters, and Remembering Written Numbers. Outcome Measures: Montreal Cognitive Assessment (MoCA overall, attention, memory), Medication-box sorting task.</p>	<p>more MoCA improvement than those without ($p<0.01$). 4. Age, education level, or type of ABI diagnosis did not have any significant effects on MoCA or medication-box scores.</p>
<p>Li et al. (2013) USA Pre-Post N=11</p>	<p>Population: ABI; Mean Age=49.45yr; Mean Time Post Injury=21.27yr. Intervention: All participants completed eight 60-min sessions using the attention and memory sub programs of the computer-based cognitive retraining Parrot Software. The participants focused on one of the eight subprograms during each session with each subprogram containing 10 lessons with increasing difficulty. Assessments were conducted before and after intervention. Outcome Measure: The cognitive assessment (attention & memory).</p>	<p>1. There was a significant improvement in attention cognitive assessment scores from pre to post intervention (mean change=2.091; $p<0.005$). 2. There was a significant improvement in memory cognitive assessment score from pre to post intervention (mean change=1.73; $p<0.05$).</p>
<p>Zickefoose et al. (2013) USA Pre-Post N=4</p>	<p>Population: TBI; Mean Age=42.75yr; Gender: Male=4, Female=0; Mean Time Post Injury=17.5yr; Severity: Severe=4, Moderate=0. Intervention: Participants engaged in computer-based brain games over the course of two 1-month treatment phases. Participants received Attention Process Training-3 (APT-3) or Lumosity™ in phase 1, and then received the alternate treatment in phase 2. Both phases consisted of twenty 30-minute sessions. Outcomes were assessed at baseline and after each phase. Outcome Measures: Test of Everyday Attention (TEA); Neurological Assessment Battery (NAB)– Numbers and Letters Test Parts B, C, and D; Perceptual rating scale (PRS).</p>	<p>1. All four participants demonstrated significant progress in reaching new levels of difficulty on all tasks over the course of both treatments ($p<0.01$). 2. NAB analysis showed that one participant demonstrated significant improvement on one sub-test, while two participants demonstrated non-significant improvement on one or more sub-tests. Improvements occurred during phase 1, regardless of treatment. 3. TEA analysis showed that one participant demonstrated improvement on several sub-tests during both treatments, while the scores of the other three participants were inconsistent for either treatment. 4. On the PRS, two participants showed strong enjoyment and willingness to continue APT-3, while the other two participants showed an equally strong rejection of APT-3 5. On the PRS, all four participants showed strong enjoyment of Lumosity™, while only two participants showed a strong willingness to continue.</p>
<p>Chen et al. (1997) USA Case-Control N=40</p>	<p>Population: Age=18+years; Gender: male=27, female=13; Condition: TBI. Intervention: Divided retrospectively into computer-assisted rehabilitation (CACR) and tradition therapy groups Outcome Measures: Neurophysiological test scores (WAIS-R; WMS).</p>	<p>1. Both groups made significant post-treatment gains on the neurophysiological test scores ($p<0.05$), with the CACR group making significant gains on 15 measures ($p<0.05$) and the comparison group making significant gains on seven measures ($p<0.005$). 2. However, no significant difference was found between groups on their post-treatment gains.</p>
	<p>Population: Cerebral infarction=23%, TBI=21%, Infection=19%, Intracerebral hemorrhage=13%, Subarachnoid hemorrhage=10%, Brain tumor=8%, Other=6%; Mean Age=43.7yr; Gender: Male=30, Female=18; Mean Time Post</p>	<p>1. At 20wk post-training, there were significant improvements in PASAT ($p<0.001$), Listening Span ($p<0.001$), Forward block repetition ($p<0.001$), Backward block repetition ($p<0.001$), COPM performance ($p<0.001$),</p>

<p>Hellgren et al. (2015) Sweden Case Series N=48</p>	<p>Injury=51.2mo. Intervention: Participants received a computerized working memory training program (Cogmed) consisting of various visuospatial and verbal working memory tasks. There were 4-5 sessions/wk for 5-7wk, consisting of 45-60min of intense exercise with one break. Occupational therapist coaches were present during every session and provided weekly feedback in addition to continuous feedback from the computer program. Outcome Measures: Paced Auditory Serial Attention Test (PASAT 2.4), Forward and backward block repetition, Listening Span Task, Canadian Occupational Performance Measure (COPM performance and satisfaction), EuroQol descriptive (EQ-5D Index), EuroQol visual analogue scale (EQ-VAS), Working Memory Index (WM Index).</p>	<p>COPM satisfaction ($p<0.001$), EQ-5D index ($p=0.009$), and EQ-VAS ($p<0.001$) compared to baseline.</p> <ol style="list-style-type: none"> 2. Compared to baseline, all participants significantly improved their WM Index at 20wk follow-up ($p<0.001$). 3. No significant differences in treatment effect were found for all outcomes in terms of sex or time post-injury, except for ≤ 18 mo since injury exhibiting more improvement than >18mo in terms of WM index difference ($p<0.05$), COPM performance improvement ($p<0.05$), and COPM satisfaction improvement ($p<0.05$).
---	---	---

Discussion

An RCT by Dirette et al. (1999) found no significant differences in improvements between participants taught specific compensatory strategies and those that completed the computer tasks without instruction of compensatory strategies. However, both groups significantly improved over time, with participants who used the compensatory strategies (whether taught or spontaneously acquired) performing better than those that did not (Dirette et al., 1999). Chen et al. (1997) also studied the effect of computer assisted cognitive rehabilitation versus traditional therapy methods. While measures of attention significantly improved in both groups after treatment, no significant differences were observed between groups (Chen et al., 1997). Similarly, an RCT by Malec et al. (1984) found no significant differences in cognitive abilities between groups of individuals who played a shooter video game.

Niemann et al. (1990) compared the effects of a computer-assisted attention retraining program with that of a memory training program, that served as a control condition. The authors found that individuals performed better on some measures of attention (Attention Test 2d) but not others (PASAT). Overall, the experimental group demonstrated significant improvements on measures of attention, when compared to the control group (Niemann et al., 1990). In the study by Gray et al. (1992), individuals received either computer-based attentional retraining or recreational computing. The authors only minor differences in attention at the end of the training; however, the experimental group performed better on the PASAT and the WAIS-R attention measures at the 6-month follow-up (Gray et al., 1992). Ruff et al. (1994) studied the effects of attention training and memory training. The authors found that THINKable, a computer-based multi-media program, was successful in improving attention in individuals with ABI (Ruff et al., 1994).

Studies with brand name computer-assisted cognitive rehabilitation have shown limited effects on attention. Lesniak et al. (2020) found that a three-week comprehensive cognitive training program using RehaCom software significantly improved attention, when compared to a three-week waiting list control condition. In a case series study, Hellgren et al. (2015) found that a computerized training program using CogMed software was successful in improving attentional scores on the Paced-Auditory Serial Attention Test. A small pre-post study examining the program Lumosity™ showed improvements in attention for a minority of participants; however, this improvement did not significantly differ from those who received Attention Process Training-III (Zickefoose et al., 2013). Parrot software showed mixed results in a pilot study reporting significant improvement in attention post-intervention (Li et al., 2013); however, subsequent study reported no significant changes on measures related to attention in individuals with ABI (Li et al., 2015). BrainHQ, an online computerized cognitive exercise program, did not significantly improve attention outcomes in individuals with ABI over time or compared to no intervention (O'Neil-Pirozzi & Hsu, 2016). The lack of evidence supporting the efficacy of computer-based cognitive rehabilitation may be due to different programs, strategies used to train participants and outcome measures used.

Conclusions

There is level 2 evidence that compensatory strategies may result in better performance in attention tasks during computer-based training for individuals with ABI (Dirette et al., 1999).

There is level 2 evidence that THINKable, a computer-based multi-media program is effective for improving attention in individuals with ABI (Ruff et al., 1994).

There is level 1b evidence that a computer-assisted attention retraining program may be more beneficial for individuals with ABI than memory training or recreational computing, when compared (Gray et al., 1992; Niemann et al., 1990).

There is level 1b evidence that video game play may not improve sustained attention in individuals with ABI (Malec et al., 1984).

There is level 2 evidence that rehabilitation interventions using computer programs and software have limited effects on attention in individuals with ABI (Hellgren et al., 2015; Lesniak et al., 2020; Li et al., 2015; Li et al., 2013; O'Neil-Pirozzi & Hsu, 2016; Zickefoose et al., 2013).

KEY POINTS

- Computer-based interventions may not be more effective than no intervention in improving measures of attention and concentration post ABI. However, a computer-assisted attention retraining program may improve attention in individuals with ABI, when compared to memory training or recreational computing.

Virtual Reality

Virtual reality (VR) is a computer technology that can generate three-dimensional interactive environments, and it has been used in recent years to provide cognitive rehabilitation to individuals living with TBI (Alashram et al., 2019). The use of VR is considered to be a safe and feasible intervention for individuals living with moderate to severe TBI, and can potentially facilitate for modification of tasks and levels of difficulty according to the needs of the individual while providing feedback in real time (Moraes et al., 2021).

TABLE 8 | The Effect of Computer-Based Virtual Reality Interventions on Attention and Reaction Time Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
Grealy et al. (1999) Scotland RCT PEDro=1 N=13	<p>Population: TBI; Age Range: 19-64yr; Gender: male=8, female=5.</p> <p>Intervention: Crossover design: participants were allocated to 4-week interventions of receiving a single bout of Virtual reality (VR) exercise or a no-exercise control condition.</p> <p>Outcome Measures: Tests measuring attention, information processing, learning, memory, and reaction and movement times.</p>	<ol style="list-style-type: none"> 1. Intervention group (n=13) performed significantly better than control group (n=320) on digit symbol ($p<0.01$), verbal ($p>0.01$) and visual ($p<0.05$) learning tasks. 2. Reaction ($p<0.01$) and movement ($p<0.05$) times improved significantly after a single VR session.
Dahdah et al. (2017) USA Pre-Post N _{Initial} =21 N _{Final} =15	<p>Population: CVA=6, TBI=5, Tumor=2, Anoxia brain injury=2; Mean Age=40.3yr; Gender: Male=12, Female=3.</p> <p>Intervention: Participants received the virtual reality (VR) intervention sessions (apartment and classroom) twice per week for a 4wk period. Sessions 1 and 8 included all types of distractors, sessions 2 and 3 included no distracting stimuli, sessions 4 and 5 included only auditory distracting stimuli, and sessions 6 and 7 included only visual distracting stimuli.</p> <p>Outcome Measures: Woodcock-Johnson, 3rd edition (WJ-III pair cancellation subtest), Delis-Kaplan Executive Function System (D-KEFS Color-Word Interference subtest), Automated Neuropsychological Assessment Metrics (ANAM Go/No-Go and unimodal Stroop subtests), VR Stroop task (apartment and classroom).</p>	<ol style="list-style-type: none"> 1. No statistically significant performance differences were found from baseline to conclusion of the study for the VR apartment Stroop or D-KEFS Stroop test. 2. For the VR classroom, participants' shortest response time on the word-reading condition was significantly reduced by session 8 ($p=0.0383$). All other VR classroom Stroop variables did not show significant differences. 3. No significant differences from session 1 to session 8 were found for all pair cancellation subtest scores. 4. From session 1 to 8, the ANAM Stroop word-reading percentage of items with a correct response ($p=0.0293$), ANAM Stroop word-reading number of correct responses per minute ($p=0.0321$), and ANAM Go/No-Go number of impulsive/bad responses ($p=0.0408$) significantly increased. All other ANAM variables did not show significant differences.
Gerber et al. (2014) USA Pre-Post	<p>Population: TBI; Mean Age=50.4yr; Gender: Male=11, Female=8; Mean Time Post Injury=10yr; GCS=4-14; Severity: Severe=9, Moderate=1, Mild=7.</p>	<ol style="list-style-type: none"> 1. All the participants reported a high level of engagement during the interactions.

<p>N=19</p>	<p>Intervention: Participants completed a series of virtual reality tasks in a standardized order utilizing a hepatic stylus; 1) Participants were asked to clear a workbench and mount tools on an upright pegboard (TOOL), then 2) spell as many 3-letter words as possible from a set of letter tiles (SPELL), then 3) prepare a virtual peanut butter and jelly sandwich (SAND), and finally 4) hammer in two nails and tighten two screws through tool use (TUSE). TOOL, SAND and TUSE tasks had a time limit of 5 minutes while SPELL task had a time limit of 2 minutes. Participants had 3 chances to perform each task (Baseline, 2nd, Final). Outcome Measures: Self-reported measures (engagement and frustration), Boredom Propensity Scale (BPS), Purdue Pegboard Test (PPT), and Neurobehavioural Symptom Inventory (NSI).</p>	<ol style="list-style-type: none"> 2. Thirty percent of participants reported a high level of frustration but were able to complete the tasks with short breaks. 3. From baseline to final, TOOL mean time decreased by 60s, TUSE mean time decreased by 68s, SAND mean time decreased by 72s and SPELL means increased by 2.7 words. 4. PPT correlated with TOOL ($p=0.016$) and TUSE ($p=0.014$) time during the final trial. 5. SPELL correlated with the BPS ($p=0.08$) during the baseline and NSI ($p=0.05$) during the final trial.
<p>Dvorkin et al. (2013) USA Pre-Post N=21</p>	<p>Population: TBI; Mean Age=37.8yr; Gender: Male=17, Female=4; Mean Time Post Injury=10.3wk. Intervention: Participants completed a virtual reality task and were instructed to hold the handle of a robot, moving the handle towards targets that appeared in the virtual environment. Participants reached to as many targets as they could within 4 minutes (1 block). Participants completed 6 blocks per day for 2 consecutive days. On each day, each pair of blocks included one haptic condition that affected the robotic handle and was either: 1) no haptic feedback (no force condition), 2) a break-through force, similar to popping a balloon (break-through condition) or 3) a gentle pulse of force (nudge condition). Outcome Measures: Tolerance, attention (pauses, pause duration), number of targets reached, and Agitated Behaviour Scale (ABS).</p>	<ol style="list-style-type: none"> 1. The interactive virtual environment was well tolerated by 18 of the 21 participants, 3 participants could not complete the 6 blocks in each visit due to fatigue or frustration. 2. In 15 participants ABS was reduced on the second visit. 3. Attention loss was reported before and during arm movements, however on the second visit participants exhibited significantly less pauses ($p<0.0001$) and shorter pause duration ($p=0.007$). 4. Participants were able to reach more targets on the second visit compared to the first visit ($p<0.0001$). 5. During the first visit, participants reached significantly less targets in the break-through and no force conditions compared to the nudge condition ($p<0.02$); the break-through and no force conditions were not significantly different. 6. During the second visit, participants reached significantly more targets in the nudge and no force conditions compared to the break-through condition ($p<0.002$); the nudge and no force conditions were not significantly different. 7. Break-through trials were significantly longer than the no force and nudge conditions on both the first and second day ($p<0.0001$). 8. Participants acquired more targets during the second visit compared to the first ($p=0.0003$) and acquired more targets with each block ($p<0.0001$).

Discussion

Repetition of tasks in VR improved attention performance over baseline for individuals with TBI (Dvorkin et al., 2013; Gerber et al., 2014). Gentle nudges corrected behaviour better than break-through or no feedback, and gentle nudges were beneficial for participants during the task. Dahdah (2017) found that individuals living with brain injury showed improved sustained attention and attention to visual details during a VR intervention that resembled the person's home environment; additionally, participants showed improvements in processing speed across sessions of VR that resembled work or school environments. However, there were no strong statistical differences (Dahdah et al., 2017). A 4-week VR exercise program in a virtual environment demonstrated significant benefits in reaction and movement times, as well as consistency and concentration, but not attentional processes after intervention (Grealy et al., 1999).

In a systematic review, Manivannan et al. (2019) examined the effectiveness of VR interventions on neurocognitive performance in individuals with TBI, including attention measures. The authors found that VR interventions may enhance attention in individuals with TBI; additionally, VR may be particularly engaging for participants and a more enjoyable experience that might result in optimized improvement (Manivannan et al., 2019).

Conclusions

There is level 2 evidence that exercising in a virtual environment may improve concentration and reaction time but may not be effective in improving attention in individuals with ABI (Grealy et al., 1999).

There is level 4 evidence that attention performance can be improved in individuals with ABI through repetition of task in virtual reality environments (Dahdah et al., 2017; Dvorkin et al., 2013; Gerber et al., 2014).



KEY POINTS

- Repetitive tasks in a virtual reality environment are effective in improving attention and concentration in individuals with ABI.

Attention Training Programs

Cicerone et al. (2005) recommended strategy training for persons with TBI for improving deficits of attention. It should be noted; however, that there was insufficient evidence to distinguish the effectiveness of specific attention training during acute stage rehabilitation from improvements made from spontaneous recovery or from more general cognitive interventions (Cicerone et al., 2005).

TABLE 9 | The Effect of Attention Training Programs on Attention and Concentration Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
<p>Dundon et al. (2015) Ireland RCT PEDro=3 N=26</p>	<p>Population: TBI; Mean Age=38.96yr; Gender: Male=19, Female=7. Intervention: Participants were assessed during a dichotic listening task (DLT) presented at 6 levels of distraction difficulty, and randomly received either adaptive training (AT, n=9), non-adaptive training (NAT, n=8), or no training (NT, n=9) between sessions (Study 2). Outcomes were assessed before and after training. Outcome Measures: DLT performance; Test of Everyday Attention (TEA).</p>	<ol style="list-style-type: none"> 1. For the DLT, there was a significant main effect of group ($F=3.99$, $p=0.035$), such that the AT group showed poorer performance than the NAT group ($p=0.019$) and the NT group ($p=0.031$). 2. For the DLT, there was a significant interaction between group and time ($F=4.38$, $p=0.026$), such that improved performance was seen in the AT ($p=0.036$) and NAT ($p=0.0025$) groups over time, but not in the NT group ($p=0.34$). 3. On the TEA, there was a significant main effect of group ($F=2.45$, $p=0.13$), such that the NT group showed better performance than the AT group ($p<0.001$) and the NAT group ($p=0.036$). 4. On the TEA, there was a significant main effect of time ($p=0.022$), such that performance improved in all groups.
<p>Cantor et al. (2014) USA RCT PEDro=7 N=98</p>	<p>Population: TBI; Mean Age=45.3yr; Gender: Male=37, Female=61; Mean Time Post Injury=12.6yr; Severity: Mild=49, Moderate=19, Severe=30. Intervention: Participants were randomly assigned to either immediate start (IS; n=49) or waitlist control (WL; n=49) groups. Participants received group sessions of emotional regulation (2 sessions, 45min) and an individual problem-solving session of attention training (1 session, 60min) per day (3 days/wk for 12 weeks). Group sizes were generally 4-6 participants. Outcome Measures: Attention Rating and Monitoring Scale (ARMS), Behavioural Assessment of the Dysexecutive Syndrome, Difficulties in Emotion Regulation Scale (DERS), Executive Function Composite from Factor Analysis (EF index), Problem Solving Inventory (PSI), and Frontal System Behavioural Scale (FrSBe).</p>	<ol style="list-style-type: none"> 1. There was a significant treatment effect for the EF index favoring the IS group ($p=0.008$). 2. There was no significant difference between groups in the DERS of ARMS. 3. Secondary analysis revealed a significant treatment effect for the FrSBe scale ($p=0.049$) and the PSI ($p=0.016$). 4. There were no other significant treatment effects. Variance of depression, age, severity, and time since injury did not change treatment effects.
<p>Amos (2002) Australia RCT PEDro=4 N=32</p>	<p>Population: TBI=16, CVA=6, Other=2, Healthy Controls=8. <i>Experimental Group</i> (n=24): Mean Age=35.71yr; Gender: Male=17, Female=7; Mean Time Post Injury=5.96yr. <i>Control Group</i> (n=8): Mean Age=31.25yr; Gender: Male=2, Female=6. Intervention: Participants with ABI were randomized into three treatment groups: unaided (n=8), external inhibition (n=8), and increased stimulus salience (n=8). All treatment</p>	<ol style="list-style-type: none"> 1. There were no significant differences in total errors between groups ($p=0.138$), but groups differed significantly in total number of trials ($p=0.025$), perseveration ($p=0.033$) and categories achieved ($p=0.001$). 2. The unaided ABI group compared to the aided ABI group (inhibition and salience) had significantly more trials ($p<0.001$), preservative errors ($p<0.006$) and lower categories score ($p<0.001$).

	<p>groups were compared to the non-ABI controls (n=8). Outcome Measures: Wisconsin Card Sorting Test (WCST).</p>	<p>3. Comparisons between the inhibition and salience aid group revealed significance difference only for perseverative errors (p<0.045); the external inhibition group displayed much less.</p>
<p>McMillan et al. (2002) UK RCT PEDro=5 N=130</p>	<p>Population: TBI; <i>Mindfulness Meditation Technique. Attentional Control Training (ACT; n=44):</i> Mean Age=34.6yr; Gender: Male=35, Female=9; Median GCS=9. <i>Physical Exercise (PE) Group (n=38):</i> Mean Age=31.4yr; Gender: Male=30, Female=8; Median GCS=10. <i>Control Group (n=48):</i> Mean Age=36.2yr; Gender: Male=36, Female=12; Median GCS=9 Intervention: Participants were assigned to 1 of 3 groups. The ACT group received supervised practice (5, 45min session over 4wk) and were given an ACT audiotape to practice daily with. The PE group had the same amount of therapist contact, but the audiotape was based on physical training. The control group had no therapist contact. Assessments were done pre- and post-training, and 6 and 12mo. Outcome Measures: Test of Everyday Attention, Adult Memory and Information Processing Battery, Paced Auditory Serial Addition Test, Trail Making Test, Sunderland Memory Questionnaire, Cognitive Failures Questionnaire.</p>	<p>1. Results showed no significant differences in outcome measures among the 3 training groups at any of the assessment points. 2. The exception to the above finding was the results of the Cognitive Failure Questionnaire where patients in both treatment groups (ACT and PE) had significantly greater reduction in self-reported cognitive failures compared to the control group at 12mo follow-up (p<0.05). 3. ACT of this duration and intensity is not recommended for individuals with attentional problems post closed head injury.</p>
<p>Sohlberg et al. (2000) USA RCT PEDro=8 N=14</p>	<p>Population: TBI=11, ABI=1, Other=2. <i>Attention Process Training (APT) Group (n=7):</i> Mean Age=33.1yr; Mean Time Post Injury=7.5yr; <i>Control Group (n=7):</i> Mean Age=38.1yr; Mean Time Post Injury=1.6yr. Intervention: Participants were randomized to receive either the APT training (treatment) or the brain injury education and supportive listening (control), in a cross over design. APT was 24hr over 10wk and the control group received 10hr over 10wk. All individuals worked directly with a therapist and assessed pre and post intervention. Outcome Measures: Trail Making Test, Paced Auditory Serial Addition Task (PASAT), Gordon Diagnostic Vigilance and Distraction, Controlled Oral Word Association Task (COWAT), Stroop Task, Attention Questionnaire.</p>	<p>1. Those in the APT group reported significantly more changes than the control group (0.91 and 0.58 respectively, p<0.05). 2. The effect of type of change was significant (p<0.0001); a greater number of memory/attention changes were reported for the APT group, whereas more psychological changes were reported for the control. 3. Changes in PASAT scores corresponded with perceived cognitive improvement in the interview; changes in PASAT scores were greater for those who reported >2 cognitive changes (p<0.05). 4. Results of the PASAT, Stroop Task, Trail Making Test B, and COWAT also found that those with higher levels of vigilance had improved scores (p<0.01). 5. For the aforementioned tasks, there were also specific improvements in performance associated with APT that were greater than those associated with brain injury education (p<0.05).</p>
<p>Fasotti et al. (2000) Netherlands RCT</p>	<p>Population: TBI; <i>Experimental Group (n=12):</i> Mean Age=26.1yr; Gender: Male=8, Female=4; Mean Time Post Injury=9.8mo. <i>Control group (n=10):</i> Mean Age=30.1yr; Gender: Male=7,</p>	<p>1. Training improved performances in both HG and WB tasks, but differences were not significant relative to control.</p>

<p>PEDro=5 N=22</p>	<p>Female=3; Mean Time Post Injury=8.3mo. Intervention: Participants in the experimental group received Time Pressure Management (TPM) training (1hr, 2-3x/wk, 2-3wk). TPM training used videotaped short stories. The program was designed to increase awareness of errors and deficits, encourage the acceptance and acquisition of the TPM strategy, and emphasize strategy application and maintenance. The control group received concentration training (30min, 2-5hr/wk, 3-4hr). Mean training was 7.4hr and 6.9hr for the TPM and control groups, respectively. Patients were assessed 2wk prior to training, post-training, and at 6mo follow-up. Outcome Measures: Waterbed (WB) and Harvard Graphics (HG) tasks, Rey's 15-word test, Rivermead Behavioural Memory Test, Auditory Concentration Test, Paced Auditory Serial Addition Task, Visual Choice Reaction Time Task.</p>	<ol style="list-style-type: none"> Scores on 2 of 3 standardized memory variables and all 3 attention variables increased significantly in the TPM group ($p<0.05$), whereas no memory variables and 1 of 3 attention variables increased significantly for the control group. Follow-up (6 mo) data for 10 from the TPM group and 9 from the control group indicated that there was a significant time effect ($p<0.05$) but no significant group time interaction ($p=0.23$); this suggests that there still was a significant improvement after 6 mo but that this improvement could not be attributed specifically to the treatment or control training.
<p>Bosco et al. (2018) Italy Pre-post N=19</p>	<p>Population: Severe TBI: Mean age=38.5yr; Gender: Male=16, Female=3; Mean time post-injury=99.4 months; GCS<8. Intervention: Groups of 5-6 participants met twice a week for 12 weeks for a total of 24 Cognitive Pragmatic Treatment (CPT) sessions. Participants were assessed at four time points, 3-months pre-treatment, immediately before treatment, immediately following treatment, and 3-months post-treatment. Outcome Measures: Assessment Battery for Communication (ABaCo), Communications Activities of Daily Living (CADL), Aachener Aphasia test, Attentional Matrices, Trail Making test, Verbal Span, Corsi's Block-Tapping test, immediate and deferred recall test, Tower of London test, Modified Card Sorting test, Raven Colored Progressive Matrices, Sally & Ann, Strange Stories.</p>	<ol style="list-style-type: none"> There was a significant difference in scores on the ABaCO between pretreatment and posttreatment scores ($p<0.001$). There were no significant differences between the two initial time points or the two posttreatment timepoints. Similar results were seen for the CADL, with individuals showing a significant improvement in their functional communication skills following treatment ($p=0.024$). Between immediate pretreatment scores and immediate posttreatment scores significant differences were only seen on the Verbal Span ($p=0.045$), and the Modified Card Sorting test ($p=0.004$). Differences in attention, spatial memory, long-term memory, planning and ToM were not significant.
<p>Serino et al. (2007) Italy Case Series N=9</p>	<p>Population: TBI: Age range=16-57 yr; Gender: male=6, female=3; Time since injury=6-78 months. Intervention: A long sequence of numbers is presented, and patients were asked to add each new number to the number preceding it and say the sum out loud. Two additional tests (the Months tasks and the Word tasks) were also administered in a similar way. The GST and the WMT were each 4 sessions/week, for 4 weeks. To vary tasks and their level of difficulty, in the interstimulus interval was varied. Outcome Measures: Working memory training</p>	<ol style="list-style-type: none"> Study results indicate the greatest improvement in performance occurred from the intermediate to the final sessions ($p<0.0005$) after the WMT. Improvement from the initial to intermediate sessions did not show any significant improvement in working memory ($p<0.46$) after GST. Working memory ($p<0.05$), divided attention ($p<0.05$), executive function ($p<0.05$), and long-term memory ($p<0.05$) for participants were significantly improved in the final session compared to the intermediate session. The same was not noted on the speed

	(WMT); Paced Auditory Serial Addition Test (PASAT); Months task	processing and sustained attention tasks ($p>0.05$). Working memory training tasks were also found to improve scores on various psychosocial outcomes.
Boman et al. (2004) Sweden Pre-Post N=10	<p>Population: TBI: Mean age=47.5yr; Gender: male=5, female=5; Time Post injury=9-40 months.</p> <p>Intervention: Each participated in an individual cognitive training session for 1 hr/3x a week for 3 weeks at home or work. The program included attention process training (APT), generalization for training and teaching of compensatory strategies for self-selected cognitive problems. Identification of cognitive problems in everyday life was also part of the compensatory strategy.</p> <p>Outcome Measures: Digit Span Test; Claeson-Dahl test; Rivermead Behavioural Memory test (RBMT); Assessment of Motor and Process Skills; European Brain Injury Questionnaire.</p>	<ol style="list-style-type: none"> 1. For the following: sustained attention, selective attention, and alternating attention significant changes ($p<0.05$, $P<0.05$, $p<0.01$ respectively) were noted in the scores of the APT test and Digit Span task between the pre to post training session and the 3 mo follow up. 2. Score increases ($p<0.05$) on the RBMT were found at the 3 mo follow up compared to the RBMT scores at the pretest. 3. When looking at changes in the RBMT score pre to post training, changes were not found. 4. No significant changes were found (pre to post and pre to 3 mo follow up) when looking at the scores on the Claeson-Dahl Memory.
Park (1999) Canada Case-Control N=46	<p>Population: TBI=23; Age matched controls=23.</p> <p>Intervention: Attention process training program of 20 two-hour sessions for a total of 40 hr.</p> <p>Outcome Measures: Paced Auditory Serial Addition Task (PASAT); Consonant Trigrams; Beck Depression Inventory (BDI).</p>	<ol style="list-style-type: none"> 1. No statistically significant improvements on the BDI from pre- to post-treatment for the TBI group. 2. TBI ($p<0.01$) and control ($p<0.001$) groups improved significantly in PASAT before/after tests. 3. Performance declined with increases in delay ($p<0.001$), and study position ($p<0.001$) on the Consonant trigrams.

Discussion

An RCT by Dundon et al. (2015) examined the effect of adaptive training on dichotic listening tasks and attention. Adaptive training involves tracking performance to adjust training difficulty according to the individual’s momentary capacity; therefore, ensuring that the individual remains challenged throughout the training (Dahlin et al., 2008). Dundon et al. (2015) found that the adaptive training group had significantly higher scores on the listening task compared to non-adaptive training group; however, the non-adaptive training group surpassed the adaptive training group in Test of Everyday Attention (TEA) scores. Overall, both groups significantly improved on measures of attention over time (Dundon et al., 2015).

Attention process training (APT) has demonstrated greater results in attention remediation compared to education alone (Sohlberg et al., 2000). A pre-post study by Boman et al. (2004) also found that APT for three weeks significantly improved attention task scores compared to pre-test scores. Park (1999) examined whether APT had a beneficial effect on attention measures (PASAT, Consonant Trigrams) in a group with severe TBI, tested pre and post training approximately 7 months apart. The author compared their results to a convenience sample of controls, given the same measures one week apart without training. Results suggested that the APT did not have a significantly beneficial effect as performance

improved on all measures across both groups, indicating practice effects and possibly spontaneous recovery (Park, 1999).

Fasotti et al. (2000) assessed the effectiveness of time pressure management (TPM) training compared to concentration training in individuals with slowed processing speed as a result of TBI. Though both groups showed improvements on information intake task performance, no significant differences between groups were observed even though specific time pressure management strategies were learned by the experimental group (Fasotti et al., 2000). Bosco et al. (2018) investigated the effects of Cognitive pragmatic treatment (CPT), a program involving executive function and theory of mind (ToM) that was developed to reintegrate individuals with TBI into their social environment (Gabbatore et al., 2015). This program was found to significantly improve scores on working memory, shifting and inhibition, as assessed by the verbal span and modified card sorting tests; however, differences in other cognitive domains, such as attention, were not significant (Bosco et al., 2018). Emotional regulation was also examined by Cantor et al. (2014). The authors assessed the effect of the Short-Term Executive Plus (STEP) program on measures of executive function, problem solving, emotion regulation and attention; however, this program was not found to be effective in the recovery of attention (Cantor et al., 2014).

The inconsistencies between studies may be due to a lack of standardized attention process training protocols. The lack of a consensus on the definition of certain cognitive processes appears to be reflected in the interventions used to attempt to rehabilitate these deficits. Unfortunately, this decreases the ability to compare studies on a more specific level; however, general conclusions can still be made that specific training programs which intend to increase attentional capacity are effective, to what extent they are more beneficial than other training programs needs to be addressed in the future through comparative methodologies. Only one study (Serino et al., 2007) described the specific task that was successful in improving attention. This cognitive task involved mental addition in combination with two other standardized tasks and was an effective strategy for improving attention.

Conclusions

There is level 2 evidence that adaptive training is no more effective than non-adaptive training in remediating attention in ABI populations (Dundon et al., 2015).

There is level 1b evidence that the Short-Term Executive Plus (STEP) cognitive rehabilitation program is not effective in treating attentional disorders compared to waitlist controls in ABI populations (Cantor et al., 2014).

There is level 2 evidence that providing an external inhibitor may improve attention switching in ABI populations (Amos, 2002).

There is level 1a evidence that attention process training (APT) may improve attentional functioning in individuals with ABI (Boman et al., 2004; Park, 1999; Sohlberg et al., 2000).

There is level 2 evidence that time pressure management training may improve attention performance and information processing speed (Fasotti et al., 2000).

There is level 4 evidence that cognitive pragmatic training (CPT) may not be effective for improving attention in individuals with ABI (Bosco et al., 2018).

There is level 4 evidence that a working memory training program may remediate attention in individuals post ABI (Serino et al., 2007).



KEY POINTS

- Attention process training may improve attentional functioning in individuals with ABI.
- Time pressure management may improve attention performance and information processing speed.

Goal Management Training

With regard to cognitive rehabilitation, therapy is typically patient-directed and driven by both long- and short-term goals (Carswell et al., 2004). The ability to self-direct towards goals is emphasized as a component of brain injury community reintegration programs and is integral in the completion of instrumental activities of daily living. The execution of these goals relies on an individual having the ability to focus attention on a given task.

TABLE 10 | The Effect of Goal Management Training on Attention and Concentration Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
<p>Chen et al. (2011) USA RCT PEDro=5 N=12</p>	<p>Population: TBI=9, Other=3: Mean Age=48yr; Gender: Male=5, Female=7; Time Post-Injury Range=6mo-6yr. Intervention: Participants were randomized to receive either the goals training intervention (n=7) or education intervention (n=5) for 5wk, after which they switched to the other condition for another 5 wk. The goals training was spread over 5wk and involved: group, individual and home-based training. The education program was a 5wk didactic educational instruction regarding brain injury. Outcome Measures: Letter number sequencing, Wechsler Adult Intelligence Scale-III, Auditory consonant trigrams, Digit Vigilance Test, Design and Verbal Fluency Switching,</p>	<ol style="list-style-type: none"> 1. In the domain of attention and executive functions, all participants in the goal training intervention showed an increase from pre to post goals training; while only 7/12 in the education intervention showed an increase from pre to post education (p<0.0001). 2. For learning and memory performance scores increased an average of 0.70 units after participation in goals training than after participation in education intervention (p=0.02). 11/12 participants improved in the goals training group while 4/12 improved in the education group (p=0.009). 3. Tests of motor speed of processing showed no significant differences between the two interventions with a non-significant trend for

	<p>Trails B, Stroop Inhibition, Hopkins Verbal Learning Test, Brief Visual Memory Test Revised, Trails A test, Visual Attention Task.</p>	<p>greater improvements in goal-training compared to education (p=0.07).</p>
<p>Novakovic-Agopian et al. (2011) USA RCT Crossover PEDro=5 N=16</p>	<p>Population: TBI=11, Stroke=3, Other=2: Mean Age=50.4yr; Gender: Male=7, Female=9; Time Post Injury Range=1-23yr. Intervention: Participants were randomized to 5wk interventions consisting of a goals training program (n=8) or an educational instruction group (n=8). Goal training focused on mindfulness-based attentional regulation and goal management strategies for participant-defined goals. Educational training was didactic instructional sessions about brain injury. At the end of 5wk, participants were switched to the other intervention. All participants were assessed at baseline, Week 5 and again at Week 10. Outcome Measures: Auditory Consonant Trigrams, Letter Number Sequencing (working memory); Digit Vigilance Test (sustained attention); Stroop Inhibition Delis-Kaplan Executive Function System (Inhibition); Trails B, Design Fluency-switching (mental flexibility), Hopkins Verbal Learning Test-Revised, Brief Visual Memory Test-Revised.</p>	<p>1. At the end of wk 5 participants in the goals-edu group showed significant improvement on measures of attention and executive function from baseline (p<0.0001), while the edu-goals group showed no change or minimal change (p>0.05). 2. The goals-edu group had significantly greater improvements than the edu-goals group on the following at wk 5: working memory (<i>Mean</i> 1.12 vs -0.12, p<0.0001); mental flexibility (<i>Mean</i> 0.64 vs 0.04, p=0.009); inhibition (<i>Mean</i> 0.62 vs 0.04, p=0.005); sustained attention (<i>Mean</i> 0.96 vs 0.27, p=0.01); learning (<i>Mean</i>=0.51 vs 0.08, p=0.02); and delayed recall (<i>Mean</i> 0.39 vs -0.27, p=0.01). 3. At wk 10, the edu-goals group significantly improved compared to wk 5 on: attention and executive function (0.79 vs 0.03, p<0.0001); working memory (1.31 vs -0.12, p<0.0008); mental flexibility (0.66 vs 0.04, p<0.0008); inhibition (0.50 vs 0.04, p=0.01); sustained attention (0.44 vs 0.27, p=0.01); memory (0.609 vs -0.10, p=0.02); learning (0.66 vs 0.08, p=0.05); and delayed recall (0.55 vs -0.27, p=0.02). 4. Those in the goals-edu group who had completed the training session were able to maintain their gains and there were significant improvements in attention and executive function (p<0.04) and working memory (p<0.02).</p>
<p>Levine et al. (2000) Canada UK RCT PEDro=4 N=30</p>	<p>Population: TBI: <i>Goal Management Training (GMT) Group</i> (n=15): Mean Age=29.0yr; Gender: Male=5, Female=10; Mean GCS=10.7; Mean Time Post Injury=3.7yr. <i>Motor Skill Training (MST) Group</i> (n=15): Mean Age=30.8yr; Gender: Male=9, Female=6; Mean GCS=10.8; Mean Time Post Injury=3.8yr. Intervention: Patients were randomized into the GMT or MST group. The GMT was comprised of five steps: 1) orienting and alerting to task, 2) goal selection, 3) partitioning goals into sub-goals, 4) encoding and retention of sub-goals, and 5) monitoring. The MST was training that was unrelated to goal management: reading and tracing mirror-reversed text and designs. Participants were tested on everyday paper and pencil tasks that focused on holding goals in mind, sub-goal analysis and monitoring. Outcome Measures: Goal Neglect (Everyday</p>	<p>Everyday paper and pencil Task 1. The GMT group compared to the MST group had significantly greater accuracy on the everyday paper and pencil tasks post-training (p<0.05). 2. The MST group also had significantly more errors during the everyday paper and pencil tasks (p<0.01). 3. The GMT group significantly reduced their errors from pre-post training during the everyday paper and pencil tasks (p<0.01). 4. The GMT also devoted significantly more time to proofreading and the room-layout tasks than the MST group from pre- to post-training (p<0.05). Neuropsychological Tasks 1. The GMT group was generally slower on timed neuropsychological tests: Stroop Interference Procedure, Trail Making Part A and B (p<0.05 and p<0.06, respectively).</p>

paper and pencil tasks), Stroop Interference Procedure, Trail Making A and B, Wechsler Adult Intelligence Scale Revised (WAIS-R).

2. No significant differences between groups for the WAIS-R ($p>0.05$).

Discussion

Levine et al. (2000) completed an RCT comparing participants using goal management training strategies to a control group exposed to only motor skills training. The treatment group improved on paper and pencil everyday tasks as well as meal preparation, which the authors used as an example of a task heavily reliant on self-regulation. Novakovic-Agonian et al. (2011), found similar results in an RCT crossover where participants were assigned to received goal-training followed by education or the reverse. The goal training first group saw a significant improvement in sustained attention compared to the education-first group, additionally the goal training first group maintained their gains over 10 weeks. A study by Chen et al. (2011) demonstrated that goal management training, although beneficial, may not be more beneficial than other interventions such as educational training with respect to processing speed. In this study both groups significantly improved in attention directed goal completion.

Conclusions

There is level 2 evidence to suggest goal management training, when compared to education, may be effective at improving attention in individuals post ABI (Chen et al., 2011; Novakovic-Agonian et al., 2011).

There is level 2 evidence that goal management training is more effective in remediating task completion times than motor skill training; however, it is not more effective in treating attention deficits, in individuals post ABI (Levine et al., 2000).



KEY POINTS

- Goal management training is effective in assisting individuals who sustained an ABI in learning to manage life goals through improved attention

Mindfulness Training

Mindfulness can be defined as a state of awareness of the present moment that is non judgemental; mindfulness interventions have the potential to improve well-being and alleviate distress, as well as to be a fruitful strategy for cognitive training (Kang et al., 2013). Mindfulness-based interventions have been used to assist neuropsychological rehabilitation in individuals with conditions such as attention deficit hyperactivity disorder (ADHD), epilepsy, multiple sclerosis (MD), Parkinson's disease (PD), and ABI (Smart et al., 2022).

TABLE 11 | The Effect of Mindfulness Training Programs on Attention and Concentration Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
McHugh & Wood (2013) Ireland RCT PEDro=5 N=24	<p>Population: TBI. <i>Mindfulness Group (N=12):</i> Mean Age=28.45yr; Mean Time Post Injury=785.5d; Mean GCS=8.5. <i>Control group (N=12):</i> Mean Age=30.5yr; Mean Time Post Injury=664.7d; Mean GCS=7.42.</p> <p>Intervention: Participants were randomly assigned to the control group or mindfulness group (focused attention). The mindfulness group received instructions (mindfulness induction) prior to completing experimental tasks. Participants then completed a memory load task (remembering the location of symbols) and an over-selectivity task and test.</p> <p>Outcome Measures: Minimal Attention Awareness Scale (MAAS), Trail making test A and B (test of visual attention and task switching) and the Wechsler Test of Adult Intelligence.</p>	<ol style="list-style-type: none"> 1. There was a significant decrease in stimulus over-selectivity after the mindfulness training compared to the control group ($p<0.05$, $t(22)=1.74$).
McMillan et al. (2002) UK RCT PEDro=5 N=130	<p>Population: TBI; <i>Mindfulness Meditation Technique. Attentional Control Training (ACT; n=44):</i> Mean Age=34.6yr; Gender: Male=35, Female=9; Median GCS=9. <i>Physical Exercise (PE) Group (n=38):</i> Mean Age=31.4yr; Gender: Male=30, Female=8; Median GCS=10. <i>Control Group (n=48):</i> Mean Age=36.2yr; Gender: Male=36, Female=12; Median GCS=9</p> <p>Intervention: Participants were assigned to 1 of 3 groups. The ACT group received supervised practice (5, 45min session over 4wk) and were given an ACT audiotape to practice daily with. The PE group had the same amount of therapist contact, but the audiotape was based on physical training. The control group had no therapist contact. Assessments were done pre- and post-training, and 6 and 12mo.</p> <p>Outcome Measures: Test of Everyday Attention, Adult Memory and Information Processing Battery, Paced Auditory Serial Addition Test, Trail Making Test, Sunderland Memory Questionnaire, Cognitive Failures Questionnaire.</p>	<ol style="list-style-type: none"> 1. Results showed no significant differences in outcome measures among the 3 training groups at any of the assessment points. 2. The exception to the above finding was the results of the Cognitive Failure Questionnaire where patients in both treatment groups (ACT and PE) had significantly greater reduction in self-reported cognitive failures compared to the control group at 12mo follow-up ($p<0.05$). 3. ACT of this duration and intensity is not recommended for individuals with attentional problems post closed head injury.

Discussion

A study by McMillan et al. (2002) examined the impact of attentional control training using mindfulness meditation on attention challenges in individuals with TBI. The authors did not find statistically significant differences on in their three treatment groups on pre-treatment measures, post-treatment or at six- or twelve-months follow-up (McMillan et al., 2002). An RCT study by McHugh and Wood (2013) found that mindfulness focused training significantly improved participants’ ability to correctly select stimuli compared to controls.

Conclusions

There is level 2 evidence that mindfulness training compared to no intervention may improve an individual’s ability to correctly reject inappropriate stimuli post ABI (McHugh & Wood, 2013).

There is level 2 evidence that an attentional control training intervention involving mindfulness meditation may not be effective to treat attentional difficulties in individuals with TBI (McMillan et al., 2002).



KEY POINTS

- Mindfulness training may offer benefits for individuals with TBI in terms of attention; however, the length of exposure needs to be carefully considered.

Music Therapy

Music therapy interventions for individuals with ABI involve the use of music (e.g., playing musical instruments, listening to music, singing) as a therapy aid to stimulate brain functions, including movement, cognition, speech, emotions, and sensory perceptions (Magee et al., 2017).

TABLE 12 | The Effect of Music Therapy on Attention and Concentration Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Martínez-Molina et al. (2021) Finland RCT Crossover PEDro=7 N _{initial} =40 N _{final} =23	<p>Population: TBI; Moderate-Severe=23; Mean Age=41.4yr; Gender: Male=10, Female=13; Mean Time Post Injury=7.8mo; Mean GCS=12.9.</p> <p>Intervention: Participants were randomized to receive usual care and neuromuscular music therapy (NMT; 60min session, twice per week for 10wk) or usual care only. NMT consisted of rhythmical training, structured cognitive motor training, and assisted music playing. Participants received individual 60min NMT sessions twice per week for a 10wk period. Outcome measures were assessed at baseline, 3mo and 6mo.</p>	<ol style="list-style-type: none"> 1. In the pre- versus post-intervention comparison, there was a significant increase (p<0.05) in temporal coupling with the sensorimotor, dorsal attention, and frontoparietal after the music-based intervention. 2. NMT increased the coupling between the frontoparietal network and dorsal attention, as well as between primary sensory networks (p<0.05).

Author Year Country Study Design Sample Size	Methods	Outcome
	<p>Outcome Measures: Frontal Assessment Battery (FAB), Wechsler Adult Intelligence Scale IV (WAIS-IV), Wechsler-Memory Scale III (WMS-III), Number-Letter Task (NLT), Auditory N-back task, Simon Task, Sustained Attention to Response Task (SART), Behavioral Rating Inventory of Executive Function-Adult version (BRIEF-A), Structural MRI (sMRI), Resting-state Functional MRI (rs-fMRI).</p>	<p>3. Decreased BRIEF-A self-monitor scores correlated significantly with decreased within-network functional connectivity in the left and right lateral prefrontal cortex nodes of the frontoparietal network (p=0.039) and showed a marginal trend with increased FAB scores (p=0.040).</p> <p>4. Participants who showed a larger reduction in connectivity within the frontoparietal network after training exhibited greater improvement in general enhanced function.</p>

Discussion

In an RCT, Martinez-Molina et al. (2021) analyzed resting-state fMRI data to assess networks associated with cognition impairment in individuals with moderate to severe TBI. The neurological therapy session involved 20 individual therapy sessions with a trained music therapist over a period of 3 months. The session targeted rehabilitation of executive function, attention and working memory. In the session, individuals participated in rhythmical training, structured cognitive motor training and assisted music playing. The authors found that neurological music therapy can lead to functional neuroplasticity changes in resting- state networks after a TBI. This study showed that music therapy can be an effective rehabilitative tool for individuals with TBI (Martinez-Molina et al., 2021).

Conclusions

There is level 1a evidence that neuromuscular music therapy may lead to changes in functional neuroplasticity, enhancing cognitive function, such as attention, in a population with moderate to severe TBI; however, further studies are needed (Martinez-Molina et al., 2021).



KEY POINTS

- Music therapy may improve cognitive function, such as attention, in individuals with severe to moderate TBI but more studies are needed.

Dance Therapy

Dance is a performing art that contributes to the integration of mind and body. Dance has been used as therapy for individuals living with TBI and may assist in the development of mind-body awareness (Winters Fisher, 2019).

TABLE 13 | The Effect of Dance Therapy on Attention Post TBI

Author Year Country Study Design Sample Size	Methods	Outcome
<p>Sarkamo et al. (2021) Finland RCT Crossover PEDro=7 N=11</p>	<p>Population: TBI: Moderate-Severe; AB Group (n=6): Mean Age=36.3yr; Gender: Male=3, Female=3; Mean Time Post Injury=9.2yr. BA Group (n=5): Mean Age=33.7yr; Gender: Male=4, Female=1; Mean Time Post Injury=5.8yr. Intervention: Participants with TBI were randomized into two groups and received the dance intervention either during the first 3-month phase (AB group) or the second 3-month phase (BA group). The Dual-Assisted Dance Rehabilitation (DARE) featured a combination of dance training and specialized physical therapy, and was provided for 60 min per day, 2 days a week, for 12 weeks. Outcome measures were assessed at the 3- and 6-month stages. Outcome Measures: Montreal Cognitive Assessment (MoCA), Frontal Assessment Battery (FAB), Wechsler Adult Intelligence Scale IV (WAIS-IV), Sustained Attention to Response Test (SART), Beck Depression Inventory II (BDI-II).</p>	<ol style="list-style-type: none"> 91% of participants were fully consistent with the protocol, and 83-100% of participants self-adhered to DARE sessions. There was a higher than-average benefit for two domains of the questionnaire on self-reported benefits of DARE: Mobility (p=0.013) and Cognition (p=0.032), as well as for the overall benefit score (p<0.001). The most consistent positive, medium-large effect sizes favouring DARE were observed for the Digit Span (p=0.232), Similarities (p=0.005), and Block Design (p=0.297) subtests of the WAIS-IV, indicating improvement in verbal working memory and reasoning ability. Both groups yielded a significant positive change from baseline to post-intervention in WAIS-IV (p=0.005) and BDI-II (p=0.002).

Discussion

The effectiveness of dance therapy for individuals with severe TBI has been observed previously in case studies (Kullberg-Turtiainen et al., 2019). A RCT by Sarkamo et al. (2021) examined the feasibility and the effects of a dance-based intervention for individuals living with severe TBI. The intervention was demonstrated to be feasible and acceptable. Sessions involved participants directing their attention to their body parts, mentally scanning their body, and following a dance choreography with music, as well as stretching and relaxing at the end of the session. Participants showed meaningful gains in motor control and mobility, as well as in memory; however, attention outcomes did not show statistically significant improvements (Sarkamo et al., 2021).

Conclusions

There is level 1a evidence that a dance program may not improve attention in individuals with moderate to severe TBI (Sarkamo et al., 2021).



KEY POINTS

Dance therapy may not improve attention in individuals with severe TBI.

Animal Assisted Therapy

Human-animal interaction has multiple benefits for individuals, with or without special medical or mental health conditions, including social attention and behavior, interpersonal interactions, mood, stress-related parameters, self-reported anxiety and physical and mental health (Beetz et al., 2012). Animal-assisted therapy (AAT) is a therapy form that has been used to address psychosocial skills and socioemotional functioning in individuals with a variety of conditions, including individuals with stroke, aphasia, severe disorders of consciousness and brain injury (Hediger et al., 2019).

TABLE 14 | The Effect of Animal Assisted Therapy on Attention Post ABI

Author, Year Country Study Design Sample Size	Methods	Outcome
<p>Gocheva et al. (2018) Switzerland RCT Crossover PEDro=7 N=19</p>	<p>Population: Non-traumatic etiology (N=13), traumatic etiology (N=9). Intervention: All participants received both conditions in randomized order, each condition consisted of 12 sessions. The experimental condition consisted of speech, occupational or physical therapy sessions accompanied by a therapeutic animal, while the control condition consisted of the same rehabilitation interventions and did not include a therapeutic animal. All conditions were completed within 6 weeks. Outcome Measures: Attention span, alertness, instances of distraction, and concentration (all outcomes were measured through behavioral analysis).</p>	<ol style="list-style-type: none"> 1. Attention span did not differ significantly between experimental and control sessions. 2. When in the animal therapy sessions individuals displayed significantly more instances of distraction compared to control sessions ($p=0.001$). Physiotherapy sessions were significantly more effected by distractions when animals were present ($p=0.016$). Further analysis demonstrated that those with higher initial FIM scores had significantly decreased instances of distraction in animal therapy sessions ($p=0.003$). 3. During animal therapy sessions, self-assessed alertness was significantly higher ($p<0.001$). There was also a significant main effect of therapy, with higher alertness in speech therapy sessions overall ($p=0.012$). Alertness was also significantly higher in the animal therapy session when individuals had higher initial FIM scores, than those that did not in animal sessions ($p<0.001$). 4. Individuals had significantly higher rates of self-reported concentration during animal therapy sessions ($p=0.014$). Concentration was also seen to be significantly higher in speech therapy sessions regardless of animal presence ($p=0.027$), with therapy type overall having a significant effect ($p<0.001$), but no significant interaction effect. Individuals with higher initial FIM scores demonstrated higher concentration scores in sessions when animals were present compared to those who had lower initial FIM scores ($p<0.001$).

Discussion

A study by Gocheva et al. (2018) examined whether the effect of therapy in the presence of an animal (e.g., rabbit, goat, guinea pig, horse) could enhance attention span and concentration in individuals with ABI. Both the animal therapy and non-animal therapy groups produced significant improvement on measures of attention and concentration; however, the animal therapy group had a significantly larger increase in concentration (Gocheva et al., 2018).

Conclusion

There is level 1b evidence that animal assisted therapy may enhance attention span and concentration in individuals with an ABI (Gocheva et al., 2018).



KEY POINTS

- Animal Assisted Therapy may improve attention span and concentration in individuals with ABI.

Brain Stimulation Techniques

Transcranial Direct Current Stimulation

Transcranial Direct Current Stimulation (tDCS) is a technique that painlessly delivers electrical currents to specific regions of the brain. These electrical currents modulate neuronal activity through electrodes placed over the head at different regions. Three recent studies have examined the effects of tDCS on attention post ABI.

TABLE 15 | The Effect of Transcranial Direct Current Stimulation on Attention Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Rushby et al. (2021) Australia RCT PEDro=6 N=30	<p>Population: TBI; Severity: Severe; Mean Age=50.0yr; Gender: Male=21, Female=9; Mean Time Post Injury=13.90yr.</p> <p>Intervention: Participants were randomized to receive a single session of anodal transcranial direct current stimulation (tDCS, 2mA for 20min) or sham tDCS (2mA for 30s). Five minutes into tDCS, participants began N-back memory tasks, where participants press a key when presented stimulus is identical to the stimulus N back in the sequence.</p> <p>Outcome Measures: Hospital Anxiety and Depression Scale (HADS), Profile of Mood State (POMS), N-back memory task, self-reported alertness, and fatigue.</p>	<ol style="list-style-type: none"> 1. There were no statistical differences in HADS scores were reported between the active and sham tDCS sessions. 2. There was a significant time effect for the POMS-Depression scale which decreased at the completion of the session in both groups (p=0.029). 3. Participants in the active tDCS group were both slower (p=0.044) and had more variable reaction time (p=0.026) than those in the sham group. 4. Arousal during the N-back tasks was higher in the sham than active group for both 1-back (p=0.011) and 2-back tasks (p=0.013).

Author Year Country Study Design Sample Size	Methods	Outcome
		5. Significant correlations were found between task activated arousal and reaction time for the 1-back task during the active tDCS session. These relationships were found to be insignificant in the sham condition.
Sacco et al. (2016) Italy RCT PEDro=4 N=32	<p>Population: TBI. Mean Time Post Injury=8.73yr; Severity: Severe=32, Moderate=0, Mild=0. <i>Treatment Group (TG, n=16):</i> Mean Age=37.7; Gender: Male=12, Female=4. <i>Control Group (CG, n=16):</i> Mean Age=35.2; Gender: Male=14, Female=2.</p> <p>Intervention: Participants were randomized to receive transcranial direct current stimulation (tDCS, TG) or sham tDCS (CG) with computer-assisted training exercises (2/d, 5d). Outcomes were assessed at baseline (T0), before treatment (T1), after treatment (T2), and 1-month follow-up (T3).</p> <p>Outcome Measures: Test for the Examination of Attention, Divided Attention substest (DA); Repeatable Battery for the Assessment of Neurological Status (RBANS).</p>	1. For DA, the TG performed significantly better at T2 compared to T0 and T1, with faster reaction times (p=0.004) and fewer omission errors (p<0.0001). 2. For DA, the CG did not perform better at T2 compared to T0 and T1. 3. For DA, there was a significant interaction between time (T0/T1 vs T2) and group (TG vs CG), for both reaction time (p=0.05) and omission errors (p=0.03). 4. On RBANS, the TG showed a non-significant improvement in performance on attention task (p=0.057), but no improvement on visual-spatial abilities, semantic fluency, working memory, and long-term memory.
Boissonnault et al. (2021) Canada Pre-Post N=6	<p>Population: TBI; Mild=1, Moderate=2, Severe=3; Mean Age=58.3yr; Gender: Male=5, Female=1; Mean Time Post Injury=11.7wk.</p> <p>Intervention: Participants received transcranial direct current stimulation (tDCS) at an intensity of 2 mA. tDCS was administered for 20-min, three times a week for three weeks. Outcome assessments were performed before and after the intervention.</p> <p>Outcome Measures: Test of Everyday Attention (TEA), Conners' Continuous Performance Test 3rd edition (CPT-3), Delis- Kaplan Executive Function System (D-KEFS), Satisfaction and Tolerability Questionnaire.</p>	1. There were no participants who completed all nine tDCS sessions. Two participants completed eight sessions, two completed two sessions, one completed five sessions, and one did not complete any sessions. 2. Three participants completed the satisfaction questionnaire. All individuals felt overall somewhat or very satisfied with their participation and would possibly or certainly recommend the project to someone else.

Discussion

Two RCTs and one pre-post study have examined the use of tDCS to improve attention following an ABI. Sacco et al. (2016) examined the effects of tDCS on attention. Individuals in the experimental group performed better in the computer-assisted divided attention task after treatment, and improvements were maintained at 1-month follow-up evaluation. The authors found that the addition of tDCS to computer-assisted training was superior to sham stimulation for improving divided attention (Sacco et al., 2016). However, more high-level studies are needed to fully examine the potential benefits of adding tDCS to traditional attentional therapies. Rushby et al. (2021) observed that the effects of tDCS are subtle and unreliable, and that there is little evidence that tDCS enhances processing speed or accuracy of working memory. It should be noted that the effects of tDCS may vary depending on the inherent ability

of each participant. Despite these challenges, tDCS may be feasible and safe for individuals with TBI (Boissonnault et al., 2021).

Conclusions

There is level 1b evidence that participants who received active tDCS presented slower and more variable reaction time than participants in the sham group. The effects of tDCS are subtle and unreliable in terms of enhancing processing speed and working memory accuracy in individuals with ABI (Rushby et al., 2021)

There is level 2 evidence that transcranial direct current stimulation when combined with computer-assisted training (compared to sham stimulation) may improve divided attention in individuals post ABI (Sacco et al., 2016).

There is level 4 evidence that tDCS may be feasible and safe for individuals with TBI (Boissonnault et al., 2021).



KEY POINTS

- Transcranial direct current stimulation (tDCS) may be effective in remediating attentional deficits when combined with computer-assisted training in ABI populations.

Transcranial Magnetic Stimulation

Transcranial Magnetic Stimulation, single pulse (TMS) and repetitive (rTMS), is used as a diagnostic tool to explore changes in cortical excitability. rTMS has relevant therapeutic application for individuals with TBI and can potentially improve some cognitive symptoms in this population (Castel-Lacanal et al., 2014), which may include attention.

TABLE 16 | The Effect of Transcranial Magnetic Stimulation on Attention Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Neville et al. (2019) Brazil RCT PEDro=9 N _{Initial} =36, N _{Final} =30	<p>Population: TBI (Diffuse Axonal Injury). Experimental Group (n=17): Gender: Male=15, Female=2; Mean Age=32.62±12.8yr; Mean GCS=5.0±3.0. Control Group (n=13): Gender: Male=12, Female=1; Mean Age=29.0±10.4yr; Mean GCS=4.4±2.5.</p> <p>Interventions: Participants were randomized to receive 10 sessions of either repetitive transcranial magnetic stimulation (rTMS) or sham stimulation. Neuropsychological evaluations were performed at baseline, post treatment and at 90d post treatment.</p> <p>Outcome Measure: Trail Making Test (TMT) A & B, Controlled Oral Word Association Test, Stroop Test,</p>	<ol style="list-style-type: none"> 1. No significant group, time or group by time interactions were found for executive function, attention, memory, or motor function, with the exception of a significant effect due to time for executive function (p<0.001) 2. Between-group comparisons of performance on TMT Part B at baseline and after the 10th rTMS session did not differ between groups (p=0.680, p=0.341, respectively).

Author Year Country Study Design Sample Size	Methods	Outcome
	Five-Point Test, Digit Span Test (Forwards & Backwards), Symbol Digit Test, Hopkins Verbal Learning Test, Brief Visuospatial Memory Test, Grooved Pegboard Test.	<ol style="list-style-type: none"> 3. Within group comparisons showed a significant difference in only the sham group on the TMT-B, showing improvement in performance (p=0.023). 4. No significant differences were observed on any neuropsychological tests. 5. No serious adverse events were reported. There was a higher frequency of mild adverse events in the rTMS group than sham, but it was not significant.
<p>Lee & Kim (2018) South Korea RCT PEDro=7 N=13</p>	<p>Population: <i>Experimental Group (N=7):</i> Mean age=42.42yr; Gender: Male=5, Female=2; Mean time post-injury=3.85 months; Mean GCS=13.71. <i>Control Group (N=6):</i> Mean age=41.33yr; Gender: Male=4, Female=2; Mean time post-injury=3.88 months; Mean GCS=13.66.</p> <p>Intervention: Individuals received either rTMS or sham rTMS for 30mins 5 times a week, for 2 weeks.</p> <p>Outcomes: Montgomery-Asberg Depression Rating Scale (MADRS), Trail Making Test (TMT), Stroop Color Word Test (SCWT).</p>	<ol style="list-style-type: none"> 1. The experimental group experienced significant within group differences on the MADRS (p<0.05), TMT (p<0.05), and SCWT (p<0.05). 2. No significant within group differences were seen for the control group. 3. Following intervention, the experimental group had significantly lower scores on the MADRS (p<0.05), TMT (p<0.05), and SCWT (p<0.05). *Lower scores indicate improved performance on TMT and SCWT.

Discussion

Two RCTs examined the effects of rTMS on attention, with conflicting results. While Lee & Kim (2018) found significant positive effects on attention and depression when compared to sham controls, Neville et al. (2019) found that rTMS had no significant positive effects on attention. Further research is necessary to determine the efficacy of rTMS for attention remediation following ABI.

Conclusions

There is conflicting level 1b evidence as to whether rTMS compared to sham stimulation may improve attention following an ABI (Lee & Kim, 2018; Neville et al., 2019).



KEY POINTS

- Repeated transcranial magnetic stimulation (rTMS) needs to be further explored; current studies have reported conflicting results as to its effectiveness in remediating attentional deficits following an ABI.

Transcranial Photobiomodulation Therapy

Photobiomodulation therapy (PBMT) refers to low-level laser therapy that involves the use of an infrared light spectrum laser as a therapeutic measure to promote tissue biostimulation, tissue regeneration, and to help relief pain and inflammation. In neurorehabilitation, the use of PBMT has the potential to contribute to the treatment of TBI, spinal cord trauma and peripheral nerve regeneration (Rosso et al., 2018).

TABLE 17 | The Effect of Transcranial Photobiomodulation Therapy on Attention Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Carneiro et al. (2019) Brazil Pre-Post N _{Initial} =10, N _{Final} =10	<p>Population: TBI=10; Mean Age=37.8±10.2yr; Gender: Male=9, Female=1; Time Post Injury Range=4mo-4yr; Severity: Mild=0, Moderate=0, Severe=10.</p> <p>Intervention: Participants received 30 min of transcranial Photobiomodulation therapy (PBMT) 3 times per wk for 6wk, for a total of 18 sessions and 540min of irradiation. Outcome measures were assessed at baseline, 1wk following PBMT, and 3mo following PBMT.</p> <p>Outcome Measures: Cerebral Blood Flow (CBF), Beck Depression Inventory II (BDI-II), Beck Anxiety Inventory (BAI), Stroop Test- Version Victoria, Trail Making Test-Forms A and B (TMT A and TMTB), Symbol Digit Test, Rey Figure Test, Rey Auditory Verbal Learning Test (RAVLT), Verbal Fluency.</p>	<p>1. No significant differences were observed across all neuropsychological outcomes and time points with PBMT. However, CBF increased, with a significant improvement in left peak systolic velocity (p=0.007).</p>

Discussion

One pre-post study by Carneiro et al. (2019) examined the effects of transcranial photo biomodulation therapy (PBMT) on attention following TBI. The authors observed an improvement in cerebral blood flow; although, no significant improvements were observed on any neuropsychological outcomes, including attention. As only a single pre-post study evaluated the effects of PBMT on attention, further research is necessary to determine the efficacy of this intervention (Carneiro et al., 2019).

Conclusions

There is level 4 evidence that transcranial photo biomodulation therapy may not improve attention following an ABI (Carneiro et al., 2019).



KEY POINTS

- Transcranial photo biomodulation therapy may not be effective in improving attention in individuals with ABI.

Pharmacological Interventions

Donepezil

Donepezil is a centrally acting acetylcholinesterase inhibitor indicated to treat dementia (Traeger et al., 2020), and it was originally developed for improving cognitive function and memory in people with Alzheimer's disease (AD) (Cacabelos, 2007). Donepezil has also proven to be beneficial for individuals living with aphasia after a stroke, and it may contribute to significant improvements in processing speed, attention, memory, awareness and functional ability among individuals living with TBI (Swenson et al., 2021). Since evidence suggests that cholinergic dysfunction may contribute to persistent cognitive deficits for people after TBI, improvements in attention, memory, and other aspects of cognition related to the acetylcholine system are expected when cholinergic function is reduced (Arciniegas, 2003).

TABLE 18 | The Effect of Donepezil on Attention and Cognitive Functioning Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Zhang et al. (2004) USA RCT PEDro=7 N=18	<p>Population: TBI; <i>Group A (n=9)</i>: Mean Age=33yr; Gender: Male=6, Female=3; Mean GCS=9.3; Mean Time Post Injury=4.6mo; <i>Group B (n=9)</i>: Mean Age=31yr; Gender: Male=7, Female=2; Mean GCS=8.9; Mean Time Post Injury=3.9.</p> <p>Intervention: In a randomized crossover trial, Group A received oral donepezil for the first 10wk, followed by a washout period of 4wk, then followed by 10wk of placebo. Group B received the treatments in the opposite order. Donepezil was administered at 5mg/d for the first 2wk, and at 10mg/d for the remaining 8wk.</p> <p>Outcome Measures: Auditory (All) and Visual (VII) subtests of Wechsler Memory Scale-III (WMS-III), and the Paced Auditory Serial Addition Test (PASAT).</p>	<ol style="list-style-type: none"> At week 10, Group A achieved significantly better scores in All (95.4±4.5 versus 73.6±4.5; p=0.002), VII (93.5±3.0 versus 64.9±3.0; p<0.001), and in the PASAT (p≤0.001) compared to Group B. This increase in scores in Group A were sustained after washout and placebo treatment (week 24), leading to no significant differences in All (105.9±4.5 versus 102.4±4.5; p=0.588), VII (91.3±3.0 versus 94.9±3.0; p=0.397), and PASAT (p>0.1) compared to Group B at study end. Within-group comparisons showed that patients in both Group A and Group B improved significantly in All and VII (p<0.05), as well as in PASAT (p<0.001), after receiving donepezil.
Campbell et al. (2018) United States PCT N=129	<p>Population: <i>Donepezil Group (N=55)</i>: Mean Age=34.4yr; Gender: Male=80%, Female=20%; Mean time post injury=28.6d; Injury Severity=Moderate-severe. <i>Control Group (N=74)</i>: Mean Age=40.8yr; Gender: Male=71.6%, Female=28.4%; Mean time post injury=25.2d; Injury Severity=Moderate-severe.</p> <p>Intervention: Individuals were assigned to receive either donepezil or a placebo treatment for an average of 67.5 days. Those receiving donepezil started with a dosage of 5mg/day, increasing to 10mg/day over the course of 7-10 days. Follow-up assessments took place approximately 61.4 days after treatment.</p>	<ol style="list-style-type: none"> For both parts of the Trail Making Test (Part A and B), there was a significant effect of time (p<0.001, p<0.001) respectively. Demonstrating that both groups significantly improved over time. No other significant effects were found for the Trail Making Test. Likewise, in the DS, only a significant effect of time (p<0.001) was observed. For both the learning and memory components of the CVLT-II there was only a

Author Year Country Study Design Sample Size	Methods	Outcome
	<p>Outcome Measures: Trail Making Tests (Part A and B), Digit Span index (DS), California Verbal Learning Test-II (CVLT-II), Logical Memory II (LMII), Functional Independence Measure (FIM), Disability Rating Scale (DRS).</p>	<p>significant effect of time observed ($p<0.001$, $p<0.001$).</p> <ol style="list-style-type: none"> The LMII showed similar results with only a significant effect of time observed ($p=0.005$). For measures of disability, both the DRS and the FIM also only showed a significant effect of time for both groups respectively ($p<0.001$, $p<0.001$). Overall, there were no significant effects of treatment found beyond that of treatment as usual in individuals with moderate to severe TBI, however both groups did demonstrate significant spontaneous recovery.
<p>Khateb et al. (2005) Switzerland Pre-Post $N_{initial}=15$, $N_{final}=10$</p>	<p>Population: TBI; Mean age=43yr; Gender: Male=8, Female=7; Mean Time Post Injury=42mo. Intervention: Participants were administered donepezil 5 mg/day for 1mo, followed by 10 mg/day for 2mos. Outcome Measures: Stroop test, Trail Making Test (TMT), Rey Auditory Verbal Memory Test (RAVMT) and Test for Attentional Performance (TAP).</p>	<ol style="list-style-type: none"> 4 of 15 participants stopped due to side effects within the first week (e.g., nausea, sleep disorders, anxiety, dizziness, etc.). Changes on the neuropsychological evaluation show modest improvement, the comparison of the global score of all questionnaires before and after therapy was marginally significant ($p=0.058$). A significant improvement in executive function was only found for the Stroop Colour naming test (87.3 ± 22.9 to 79.5 ± 19.1, $p=0.03$); for learning and memory the RAVMT-learning (47.7 ± 6.9 to 53.5 ± 5.0, $p=0.05$); and for attention, the errors subsection of divided attention (5.8 ± 3.3 to 2.9 ± 2.7, $p=0.03$).

Discussion

In an RCT, Zhang et al. (2004) demonstrated that donepezil was associated with significantly more improvement in tasks of sustained attention compared to placebo. These improvements were sustained even after the washout period. Once both groups had completed donepezil treatment there were no significant differences between groups on any measures of attention. Khateb et al. (2005) found that individuals performed significantly better on measures of divided attention after donepezil treatment; however, 4 of 15 participants stopped treatment due to negative side-effects. In contrast to the positive effects found by these studies, one prospective controlled trial found no significant effects of donepezil on any measures of cognition, including attention (Campbell et al., 2018). In both the Campbell et al. (2018) and Zhang et al. (2004) studies, individuals received donepezil for approximately the same duration. Donepezil may be effective in enhancing cognition in individuals with TBI; however, long-term use outcomes are still unknown (Florentino et al., 2022).

Conclusions

There is conflicting level 1b (positive) and level 2 (negative) evidence that donepezil may improve attention compared to placebo post TBI (Campbell et al., 2018; Khateb et al., 2005; Zhang et al., 2004).



KEY POINTS

- It is unclear as to whether donepezil may improve attention in individuals with moderate to severe TBI.

Methylphenidate

Methylphenidate is a central nervous stimulant which inhibits the reuptake of dopamine and norepinephrine, resulting in increased dopaminergic activity. Methylphenidate is extensively used to treat attention deficit hyperactivity disorder (ADHD) in children and adults (Cândido et al., 2021). This medication has also been used to treat mental fatigue and to enhance cognitive functioning in individuals living with TBI (Levin et al., 2019). In terms of medication safety, no serious side effects have been observed in clinical trials (Godfrey, 2009). However, there is a small number of reports of adverse effects in individuals taking methylphenidate; therefore, appropriate dosing is imperative (Barnett & Reid, 2020).

TABLE 19 | The Effect of Methylphenidate on Attention, Concentration and Processing Speed Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Jenkins et al. (2019) UK RCT Crossover PEDro=9 N _{Initial} =46, N _{Final} =40	<p>Population: TBI=40; <i>Treatment Group (Intervention First; n=20):</i> Mean Age= 40±12yr; Gender: Male=18, Female=2; Mean Time Post Injury=67±85mo; Severity: Mean GCS=8.3±5.2.</p> <p><i>Control Group (Placebo First; n=20):</i> Mean Age=39±12yr; Gender: Male=16, Female=4; Mean Time Post Injury=67±85mo; Severity: Mean GCS=8.3±5.4.</p> <p>Intervention: Participants were randomized to receive 0.3mg/kg of methylphenidate (treatment group) twice a day for 2wk with crossover to placebo (control group) twice a day for 2wk and vice versa. Outcome measures were assessed at baseline, 2 and 4wk.</p> <p>Outcome Measures: Choice Reaction Time (CRT) Task, Single-Photon Emission Computed Tomography (SPECT), Trail Making Test (TMT), Stroop Test, People Test, Wechsler Abbreviated Scale for Intelligence (WASI), Lille Apathy Rating Scale (LARS), Visual Analogue Scale for Fatigue (VAS-F), Glasgow Outcome</p>	<ol style="list-style-type: none"> 1. No significant differences (p>0.05) were observed between groups on several measures: TMT, Stroop, People Test, WASI, FrSBe, GOSE, HADS, Cognitive Failures Questionnaire, or Rating Scale of Attentional Behaviour. 2. Using SPECT imaging, participants were divided into groups with low and normal dopamine transporter levels for analysis. 3. Participants with low dopamine transporter levels receiving methylphenidate significantly improved on several measures when compared to controls: CRT (p=0.02), LARS self-reported (p=0.03) and caregiver (p=0.02), VAS-F (p=0.007). 4. Participants with normal dopamine transporter levels receiving methylphenidate reported significantly less

Author Year Country Study Design Sample Size	Methods	Outcome
	Scale-Extended (GOSE), Hospital Anxiety and Depression Scale (HADS), Frontal Systems Behaviour Scale (FrSBe), Cognitive Failures Questionnaire, Rating Scale of Attentional Behaviour.	fatigue when compared to controls (VAS-F, p=0.03).
Dymowski et al. (2017) Australia RCT PEDro=9 N _{Initial} =11, N _{Final} =10	<p>Population: TBI. <i>Methylphenidate Group</i> (n=6): Mean Age=35yr; Gender: Male=4, Female=2; Mean Time Post Injury=366 d; Mean Worst GCS=4.83. <i>Placebo Group</i> (n=4): Mean Age=32.5yr; Gender: Male=2, Female=2; Mean Time Post Injury=183.5 d; Mean Worst GCS=4.50.</p> <p>Intervention: Participants were randomly assigned to receive either methylphenidate (0.6 mg/kg/d rounded to the nearest 5mg with maximum daily dose of 60 mg) or placebo (lactose). Outcomes relating to processing speed, complex attentional functioning, and everyday attentional behaviour were assessed at baseline, 7wk (on-drug), 8wk (off-drug), and 9mo follow-up.</p> <p>Outcome Measures: Symbol Digit Modalities Test (SDMT), Trail Making Test (TMT) A and B; Hayling (A, B, error), Digit Span (DS-Forward, Backward, Sequencing, Total), Ruff 2&7 Selective Attention Test Automatic Speed Raw Score (2&7 ASRS), Ruff 2&7 Selective Attention Test Controlled Speed Raw Score (2&7 CSRS), Simple Selective Attention Task Reaction Time (SSAT RT), Complex Selective Attention Task Reaction Time (CSAT RT), N-back 0-back RT, N-back 1-back RT, N-back 2-back RT, Rating Scale of Attentional Behaviour Significant Other (RSAB SO).</p>	<ol style="list-style-type: none"> After applying Bonferroni corrections, no significant differences between groups from baseline to 7 wk, baseline to 8 wk, or baseline to 9 mo were observed for SDMT, TMT A, TMT B, Hayling A, Hayling B, Hayling error, DS Forward, DS Backward, DS Sequencing, DS Total, 2&7 ASRS, 2&7 CSRS, SSAT RT, CSAT RT, N-back 0-back RT, N-back 1-back RT, N-back 2-back RT, or RSAB SO. Methylphenidate was not associated with improved everyday attentional behaviour or processing speed after traumatic brain injury.
Zhang & Wang (2017) China RCT PEDro=10 N _{Initial} =36, N _{Final} =33	<p>Population: TBI; Severity: mild to moderate. <i>Methylphenidate Group</i> (n=18): Mean Age=36.3 yr; Gender: Male=13, Female=5. <i>Placebo Group</i> (n=18): Mean Age=34.9yr; Gender: Male=14, Female=4.</p> <p>Intervention: Participants were randomly assigned to receive methylphenidate (flexibly titrated from 5 mg/d at the beginning, then gradually increased by 2.5 mg/d until reaching 20 mg/d) or placebo for 30 wk.</p> <p>Outcome Measures: Mental Fatigue Scale (MFS), Choice Reaction Time (CRT), Compensatory Tracking Task (CTT), Mental Arithmetic Test (MAT), Digit Symbol Substitution Test (DSST), Mini-Mental State Examination (MMSE), Beck Depression Inventory (BDI), Hamilton Rating Scale for Depression (HAMD).</p>	<ol style="list-style-type: none"> At baseline, there were no significant differences between groups in terms of demographics, MFS, CRT, CTT, MAT, DSST, MMSE, BDI, or HAMD. Post-intervention, the experimental group had significantly lower scores compared to control group for MFS (p=0.005), CRT (p<0.001), CTT (p<0.001), BDI (p=0.040), and HAMD (p=0.005). Post-intervention, the experimental group had significantly higher scores compared to control group for MAT (p=0.020), DSST (p<0.001), MMSE (p<0.001).
Willmott et al. (2013) Australia RCT PEDro=10 N=32	<p>Population: TBI; Gender: Male=21, Female=11; Mean Time Post Injury=68 d; <i>TBI Val/Val Group</i> (n=11): Mean Age=22.64 yr; Mean GCS=4.67; <i>TBI Val/Met Group</i> (n=14): Mean Age=28.57 yr; Mean GCS=5.38; <i>TBI Met/Met Group</i> (n=7): Mean Age=30.57 yr; Mean GCS=6.83.</p>	<ol style="list-style-type: none"> At baseline, there were no significant differences across various genotypes on attentional performance. Participants with TBI and Met/Met alleles performed significantly poorer on the SDMT (p<0.0005), 2 & 7 ASRS (p=0.001), 2 & 7 CSRS (p<0.0005), DC RT (p=0.005), and SI RT

Author Year Country Study Design Sample Size	Methods	Outcome
	<p>Intervention: Participants with TBI, in a crossover design, received 0.3 mg/kg methylphenidate (2 x/d) for 6 sessions in total (spanning 2 wk), alternating between treatment and placebo for every other session. Results were compared against those from healthy controls (n=40). Groups were stratified by the presence of the Val158Met gene.</p> <p>Outcome Measures: Ruff 2 & 7 Selective Attention Test – automatic (2 & 7 ASRS) and controlled (2 & 7 CSRS), Selective Attention Task, Four Choice Reaction Time Task (4CRT) – dissimilar compatible (DC) and similar incompatible (SI), Symbol Digit Modalities Test (SDMT), Letter Number Sequencing Task, Wechsler Test of Adult Reading.</p>	<p>(p=0.002), when compared to controls. Analyses with participants with TBI and Val/Val alleles showed even worse outcomes, demonstrating poorer performance on 7/8 outcome measures.</p> <p>3. Following methylphenidate treatment one significant drug and genotype interaction was seen between Met/Met carriers and performance on the SDMT (F=4.257; p=0.024), suggesting Met/Met carriers were more responsive to methylphenidate than either of the others.</p>
<p>Kim et al. (2012) USA RCT PEDro=7 N=23</p>	<p>Population: Moderate/Severe TBI; Mean Age=34.2 yr; Gender: Male=18, Female=5; Mean Time Post Injury=51.1 mo.</p> <p>Intervention: In a crossover design, participants were randomly assigned to receive 0.3 mg/kg methylphenidate followed by placebo, or the reverse and were assessed after each.</p> <p>Outcome Measures: Visual sustained attention task (VSAT), Two-back task.</p>	<p>1. Relative to placebo, both accuracy (1.62±1.03 versus 2.23±1.07; p<0.005) and mean reaction time (827.47±291.17s versus 752.03±356.87s; p<0.050) in the VSAT were improved significantly on MPH.</p> <p>2. Relative to placebo, mean reaction time (929.31±192.92s versus 835.02±136.12s; p<0.050), but not accuracy, in the two-back task was improved significantly when on MPH.</p>
<p>Willmott & Ponsford (2009) RCT PEDro=10 N=40</p>	<p>Population: TBI; Mean Age=26.93yr; Gender: Male=28, Female=12; Time since injury=68.38 d.</p> <p>Intervention: Participants received either methylphenidate (0.3 mg/kg 2 x/d, rounded to the nearest 2.5 mg) or a placebo. Participants were seen for 6 sessions across 2-week period, then crossed-over.</p> <p>Outcome Measures: Ruff 2 and 7 Selective Attention Test, Selective Attention Task, Four Choice Reaction Time Task, Sustained Attention to Response Task, Symbol Digit Modalities Test, Letter Number Sequencing Task, Wechsler Test of Adult Reading.</p>	<p>1. Methylphenidate significantly increased speed of information processing on the Symbol Digit Modalities Test (p=0.020); Ruff 2 and 7 Test-Automatic Condition (p=0.003); Simple Selective Attention Task (p=0.001); Dissimilar compatible (p=0.003), and Similar Compatible (p=0.002).</p>
<p>Kim et al. (2006) Korea RCT PEDro=6 N=18</p>	<p>Population: TBI; <i>Methylphenidate Group (n=9):</i> Mean Age=30.1yr; Gender: Male=9, Female=0; Mean Time Post Injury=1.6yr; <i>Placebo Group (n=9):</i> Mean Age=38.3yr; Gender: Male=7, Female=2; Mean Time Post Injury=3.6 yr.</p> <p>Intervention: Patients were randomly allocated to receive either 20 mg methylphenidate or the placebo. Assessments were done at baseline (T1), 2 hr post treatment (T2), and 2 d later (T3).</p> <p>Outcome Measures: Visual sustained attention task (VSAT), Two-back task.</p>	<p>1. At T1 there were no significant differences in mean reaction time or in accuracy between the two groups.</p> <p>2. For those in the treatment group, the mean reaction time of the two-back task improved significantly compared to those in the placebo group from T1 to T2 (13.74±13.22% versus 4.02±9.48%; p<0.05).</p> <p>3. No significant difference in improvement as seen with accuracy of the two-back task (p=0.07), nor with the VSAT.</p>
<p>Whyte et al. (2004)</p>	<p>Population: TBI; Mean Age=37 yr; Gender: Male=29, Female=5; GCS<12; Median Time Post Injury=3.2 yr.</p>	<p>1. Methylphenidate showed significant improvements in information processing</p>

Author Year Country Study Design Sample Size	Methods	Outcome
USA RCT PEDro=8 N=34	<p>Intervention: Participants received 0.3 mg/kg/dose methylphenidate for 3 wk, 2x/d, and placebo for 3 wk, for a total of 6 wk, with conditions alternating weekly. Washout lasted a day, after which time the groups crossed over.</p> <p>Outcome Measures: Sustained Arousal and Attention Task, Speed/Accuracy Tradeoff Task, Distraction Task, Choice Reaction Time Task, Dual Task, Sustained Attention and Response Task, Task of Everyday Attention, Inattentive Behavior Task, Cognitive Failures Questionnaire.</p>	<p>speed ($p<0.001$), work task attentiveness ($p=0.010$), and caregiver attention ratings ($p=0.010$), pre-post.</p> <ol style="list-style-type: none"> No treatment-related improvements were observed in susceptibility to distraction and divided or sustained attention. A dose of 0.3mg/kg of Methylphenidate, twice a day, may have a consistent positive effect on speed processing and caregivers rating of attentiveness.
Plenger et al. (1996) USA RCT PEDro=5 N=23	<p>Population: TBI; Gender: Male=17, Female=6; <i>Placebo Group (n=13):</i> Mean Age=26.6yr; Mean GCS=8.1; <i>Methylphenidate Group (n=10):</i> Mean Age=31.4yr; Mean GCS=9.3.</p> <p>Intervention: Participants were randomly allocated to receive either methylphenidate or placebo. Methylphenidate was administered at 0.30 mg/kg, 2x/d, for 30 d.</p> <p>Outcome Measures: Disability Rating Scale (DRS), Continuous Performance Test (CPT), 2 & 7 Test (2 & 7), Paced Auditory Serial Addition Test (PASAT), Digit Span & Attention/ Concentration from Wechsler Memory Scale-Revised (Attn/Conc from WMS-R).</p>	<ol style="list-style-type: none"> At 30 d follow-up (n=15), significant differences were obtained on DRS, suggesting better outcome for the methylphenidate group. This difference however was not seen at 90 d follow-up (n=11). Significant differences were found on the attention-concentration domain at the 30-day follow-up, as indicated by CPT, PASAT, 2 & 7, and Attn/Conc from WMS-R ($p<0.030$). The treatment group performed better in these measures compared to the placebo group.
Speech et al. (1993) USA RCT PEDro=7 N=12	<p>Population: TBI; Mean Age=27.6yr; Gender: Male=5, Female=7; Mean Time Post Injury=48.5 mo.</p> <p>Intervention: In a crossover design, participants were randomly assigned to receive 0.3 mg/kg methylphenidate, 2 x/d, for 1 wk, followed by 1wk of placebo, or receive the treatment in the reverse order.</p> <p>Outcome Measures: Gordon Diagnostic System, Digit Symbol, and Digit Span subtests of the Wechsler Adult Intelligence Scale-Revised, Stroop Interference Task, Sternberg High Speed Scanning Task, Selective Reminding Test, Serial Digit Test, Katz Adjustment Scale.</p>	<ol style="list-style-type: none"> No significant differences were found between methylphenidate and placebo condition in any of the outcome measures studied.
Gualtieri & Evans (1988) United States RCT Crossover PEDro=7 N=15	<p>Population: TBI, Mean age=24.1yr; Gender: Male=10, Female=5; Mean time post-injury=46.8mo.</p> <p>Intervention: Participants were assigned to receive three conditions in randomized order. 1) Placebo; 2) Methylphenidate (0.15mg/kg) twice daily; 3) Methylphenidate (0.30mg/kg) twice daily. Each condition was 12 days long, with 2 days washout between conditions.</p> <p>Outcome Measures: Adult Activity Scale self-administered (AAS-S), Adult Activity Scale (administrator)(AAS-O), Examiner’s Rating Scale (EXRS), Self Rating Scale (SRS), Verbal Fluency Test (VFT), Non-verbal Fluency test (NVFT).</p>	<ol style="list-style-type: none"> There was a significant improvement in AAS-S and AAS-O scores between the placebo and high-dose conditions ($p<0.05$). There was a significant difference in SRS scores between the placebo group and the high-dose condition ($p<0.05$). On the EXRS there was a significant difference between baseline and low-dose ($p=0.012$), placebo and low-dose ($p=0.025$), baseline and high-dose ($p=0.012$), with higher doses of methylphenidate having improved effects.

Author Year Country Study Design Sample Size	Methods	Outcome
		<ol style="list-style-type: none"> 4. There was a significant improvement in VFT scores between baseline and the high-dose groups ($p=0.017$). 5. There was a significant difference on NVFT scores between baseline and placebo ($p=0.008$), baseline and low-dose ($p=0.008$), baseline and high-dose ($p=0.008$), and the placebo and high-dose group ($p=0.018$), with methylphenidate improving scores.
<p>Whyte et al. (1997) United States RCT Crossover PEDro=7 N=19</p>	<p>Population: TBI; Mean age=30.7yr; Gender: Male=15, Female=4; Mean GCS=5.83. Intervention: Individuals were randomly assigned to either receive methylphenidate first or placebo, and then the reverse. Methylphenidate was given twice a day at a dose of 0.25mg/kg. Outcome Measures: Sustained arousal task, phasic arousal task, distraction task, choice reaction-time task, behavioral inattention.</p>	<ol style="list-style-type: none"> 1. There was a significant drug x performance interaction ($p<0.001$), where performance was differentially impacted by the drug on each assessment. 2. Group stratification revealed that methylphenidate was more effective for improving performance on attentional measures for younger participant than older ones ($p<0.05$). 3. There were no other significant effects.
<p>Pavlovskaysa et al. (2007) Pre-Post Israel N=6</p>	<p>Population: TBI; Age Range=18-47yr; Gender: Male=4, Female=2; GCS ≥ 8. Intervention: Participants were administered 5 to 10 mg of methylphenidate (MHP) over a 2-week period. Participants were evaluated before, during and after the administration of methylphenidate. Outcome Measures: Performance on the Visual Spatial Attention Task Analyzing Rightward and Leftward Shifts of Attention.</p>	<ol style="list-style-type: none"> 1. Prior to treatment, patients were found to have great difficulty in shifting attention between hemifields. 2. There was a significant improvement in the asymmetry with MHP ($p<0.001$). 3. The right-side performance was significantly better on average than the left side (0.77 versus 0.59; $p<0.050$). 4. Performance was significantly better for ipsilateral valid cueing ($p<0.010$) than for invalid cross-trials ($p<0.001$). 5. The difference between ipsi- and cross-cueing for left side target performance is significant for each of the stages ($p<0.001$).

Discussion

A variety of studies with different dosing regimens and durations have found positive effects of methylphenidate in individuals with TBI (Gualtieri & Evans, 1988; Whyte et al., 1997; Zhang & Wang, 2017). In an RCT, Whyte et al. (2004) indicated that speed of processing, attentiveness during individual work tasks and caregiver ratings of attention were all significantly improved with methylphenidate treatment. No treatment related improvement was seen in divided or sustained attention, or in susceptibility to distraction. Methylphenidate was also found to significantly improved attention and concentration, visuo-spatial attention, and reaction time (Kim et al., 2012; Pavlovskaya et al., 2007; Plenger et al., 1996). These findings align with Willmott and Ponsford (2009), who found that

administering methylphenidate to a group of individuals during inpatient rehabilitation significantly improved the speed of information processing.

Other studies that examined the effects of methylphenidate did not find significant differences compared to placebo on measures of attention, information processing speed, or learning (Dymowski et al., 2017; Kim et al., 2006; Speech et al., 1993). A potential explanation for these conflicting results is proposed by Willmott et al. (2013) who hypothesize that an individuals' response to methylphenidate depends on their genotype. More specifically, that individuals possessing the methionine (Met) allele at the catechol-O-methyltransferase (COMT) gene would confer greater response to methylphenidate compared to those with the valine (Val) allele. While both Met/Met and Val/Val carriers performed more poorly in various attentional tasks compared to healthy controls, Met/Met carriers did show greater improvements in strategic control in attention than Val/Val carriers. Jenkins et al. (2019) offer another possible explanation for these conflicting results, suggesting that the variability in treatment effect may be due to variability in dopaminergic damage between individuals. The authors found that only individuals with low dopamine transporter levels, demonstrated improvements in attention with methylphenidate.

A meta-analysis study (Huang et al., 2016) showed statistically significant evidence that Methylphenidate may help to improve attention in individuals (both adults and children) living with TBI, especially sustained attention. In line with this finding, a recent systematic review (Barnett & Reid, 2020) found that the use of methylphenidate in individuals living with ABI mostly improved cognitive abilities such as working memory, processing speed and attention. Methylphenidate had a beneficial effect on cognitive outcomes post-injury; however, more longitudinal studies are needed to explore the possibility of ceiling effects. Despite the lack of evidence regarding long-term use of this medication by individuals with TBI, Methylphenidate has demonstrated a positive effect on speed of information processing in this population (Ponsford et al., 2014).

Conclusions

There is conflicting level 1a evidence regarding the effectiveness of methylphenidate following brain injury for the improvement of attention in individuals post TBI (Dymowski et al., 2017; Gualtieri & Evans, 1988; Kim et al., 2012; Kim et al., 2006; Pavlovskaya et al., 2007; Plenger et al., 1996; Speech et al., 1993; Whyte et al., 2004; Whyte et al., 1997; Willmott & Ponsford, 2009; Willmott et al., 2013; Zhang & Wang, 2017).

There is level 1b evidence that individuals carrying the Met allele may be more responsive to methylphenidate than those without the Met allele when it comes to the TBI population (P. Jenkins et al., 2019).



KEY POINTS

- Studies examining methylphenidate for attention and information processing show conflicting results.
- Response to methylphenidate may depend on the presence of the Met genotype and/or dopamine transporter levels, as well as on the age of the individual.

Bromocriptine

Bromocriptine is a dopaminergic agonist which exerts its effects primarily through the binding of D₂ receptors (Whyte et al., 2008). It has been suggested that dopamine is an important neurotransmitter for prefrontal function (McDowell et al., 1998). Dopamine agonists such as bromocriptine have been used to treat individuals with TBI that present with minimally conscious state (Passler & Riggs, 2001), as well as to improve cognitive functioning in individuals with TBI (Frenette et al., 2012). Two examined the use of bromocriptine to facilitate cognitive recovery in individuals with TBI.

TABLE 20 | The Effect of Bromocriptine on Attention Post TBI

Author Year Country Study Design Sample Size	Methods	Outcome
Whyte et al. (2008) USA RCT Crossover PEDro=7 N=12	<p>Population: Moderate/ Severe TBI; Mean Age=35.75yr; Gender: Male=8, Female=4; Median Time Post Injury=3.3yr.</p> <p>Intervention: In a crossover design, participants were randomly assigned to receive bromocriptine (1.25 mg 2x/d titrated to 5mg 2x/d over a 1wk), followed by placebo or the reverse order. Each lasted 4wk with a 1wk washout period.</p> <p>Outcome Measures: Attention Tasks.</p>	<ol style="list-style-type: none"> Though some improvements were observed in certain subsets of attentional tasks (e.g., speed decline, decline in responding, test of everyday attention), they were not significant. Overall results suggest bromocriptine had little effect on attention.
McDowell et al. (1998) USA RCT PEDro=4 N=24	<p>Population: TBI; Median Age=32.5yr; Gender: Male=20, Female=4; GCS Range=3-8; Time Post injury Range=27d-300mo.</p> <p>Intervention: In a crossover design, participants were randomly assigned to receive 2.5 mg bromocriptine followed by placebo, or the reverse order.</p> <p>Outcome Measures: Dual-task paradigm (counting and digit span), Stroop Test, spatial delayed-response task, Wisconsin Card Sorting Test (WCST), reading span test, Trail Making Test (TMT), controlled oral word association test (COWAT), Control tasks.</p>	<ol style="list-style-type: none"> Following bromocriptine treatment there were significant improvements on the dual-task counting (p=0.028), dual-task digit span (p=0.016), TMT (p=0.013), Stroop Test (p=0.05), COWAT (p=0.02), and WCST (p=0.041). Bromocriptine had no significant effects on working memory (e.g., spatial delayed-response task and reading span test; p=0.978), or on control tasks (p=0.095).

Discussion

The effect of bromocriptine on cognitive function in individuals with TBI was explored in two RCTs (McDowell et al., 1998; Whyte et al., 2008). In the study by McDowell (1998), low-dose bromocriptine (2.5 mg daily) improved functioning on tests of executive control including a dual task, Trail Making Test, the Stroop test, the Wisconsin Card Sorting Test and the controlled oral word association. However, a later study by Whyte et al. (2008) found that bromocriptine had little effect on attention, and it was noted that several participants did experience moderate to severe drug effects and withdrew or were withdrawn from the study.

Although McDowell et al. (1998) demonstrated some benefits following administration of bromocriptine, there was only a single administration of bromocriptine and the dose was considerably lower than that given by Whyte et al. (2008). Spontaneous recovery may have been a factor leading to the improved abilities in individuals receiving a single dose (2.5 mg daily) of the medication; however, study results did not answer this question. Results from Whyte et al. (2008) noted that the placebo group demonstrated better, although not significant, trends in improvement on the various tasks administered.

Conclusions

There is conflicting evidence (Level 1b and Level 2) as to whether bromocriptine improves performance on attention tasks compared to placebo in individuals post TBI (McDowell et al., 1998; Whyte et al., 2008).



KEY POINTS

- The effectiveness of Bromocriptine on attention for individuals with ABI is unknown at this time, further studies are needed.

Cerebrolysin

Cerebrolysin has been demonstrated to have neuroprotective and neurotrophic effects and has been linked to increased cognitive performance in an elderly population. Cerebrolysin has shown protective effects against neurodegenerative injury or disease, and it has demonstrated positive effects on neuroplasticity (Fiani et al., 2021). As explained by Alvarez et al. (2003), “*Cerebrolysin (EBEWE Pharma, Unterach, Austria) is a peptide preparation obtained by standardized enzymatic breakdown of purified brain proteins, and comprises 25% low-molecular weight peptides and free amino acids*” (pg. 272).

TABLE 21 | The Effect of Cerebrolysin on Attention Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Poon et al. (2020) Italy RCT PEDro=9 N=40	<p>Population: TBI; Mean Age=38.1yr; Gender: Male=32, Female=8; Mean Time Post Injury<6hr; Mean GCS=9.9.</p> <p>Intervention: Participants were randomized to receive either Cerebrolysin (50 mL) or placebo for 10 days, followed by two additional treatment cycles (10 mL daily for 10 days). Outcomes were assessed at baseline, day 10, day 30, and day 90 of the study.</p> <p>Outcome Measures: Colour Trails Test (CTT), Hospital Anxiety and Depression Scale (HADS).</p>	<ol style="list-style-type: none"> 1. The largest effects of all outcome scales were related to the CTT 1 and 2 ($p=0.0223/0.0170$). For CTT test 1, the median completion time in the Cerebrolysin group (58s) was less than half of the time needed in the placebo group (164s). For CTT test 2, the mean completion time was 117s in the Cerebrolysin group and 240s in the placebo group. 2. Sensitivity analysis of the primary multidimensional outcome ensemble showed a statistically significant superiority of Cerebrolysin. 3. For HADS, a statistically significant superiority of Cerebrolysin was found with a large effect size ($p=0.0378$).
Alvarez et al. (2003) Spain Pre-Post N=20	<p>Population: TBI; Mean Age=30.1yr; Gender: Male=15, Female=5; Mean GCS=6.1; Time Post Injury Range=23-1107d.</p> <p>Intervention: Participants with TBI received a total of 20 intravenous infusions of cerebrolysin solution (30 mL/infusion) over 4wk. Assessments were made at baseline, during treatment, and after the 4wk treatment period.</p> <p>Outcome Measures: Syndrome Kurztest (SKT), Electroencephalogram (EEG)/brain mapping recordings, Glasgow Outcome Scale (GOS).</p>	<ol style="list-style-type: none"> 1. Compared to baseline, participants with TBI showed a significant decrease in slow bioelectrical activity frequencies (δ: $p<0.010$; θ: $p<0.050$), and a significant increase in fast frequencies (β: $p<0.010$) after receiving cerebrolysin, suggesting improvement in brain bioelectrical activity. 2. Significant improvements in SKT performance were noted from pre to post treatment (15.9 ± 2.4 versus 12.0 ± 2.1; $p<0.010$). 3. GOS scores significantly improved from pre to post treatment (3.7 ± 0.3 versus 3.95 ± 0.3; $p<0.050$).

Discussion

In an open-label trial of 20 participants with TBI, Alvarez et al. (2003) found that cerebrolysin was associated with improved brain bioelectrical activity, as evidenced by a significant increase in fast beta frequencies. A brief neuropsychological battery (Syndrome Kurztest) consisting of nine subtests was administered to evaluate memory and attentional functions in participants undergoing treatment with cerebrolysin. There was an overall significant improvement in performance post treatment, suggesting they experienced cognitive benefits from cerebrolysin treatment. Improvements in the Glasgow Outcome Scale were also observed (Alvarez et al., 2003). More recently, a randomized placebo-controlled multi-site trial of individuals living with TBI also demonstrated the benefits of cerebrolysin, compared to the placebo (Poon et al., 2020). The results of this study showed the relevance of multimodal agents for individuals living with TBI, due to the combination of pleiotropic neuroprotective effects with neuroregeneration enhancement. A recent meta-analysis (Vester et al., 2021) also

confirmed that Cerebrolysin is a safe and effective treatment for individuals living with moderate to severe TBI. Together these findings suggest that cerebrolysin may represent an effective neuroprotective therapy with tangible cognitive benefits for individuals living with an ABI. However, additional controlled trials are necessary to further explore the efficacy of this drug.

Conclusion

There is level 1b evidence that Cerebrolysin is an effective therapy that may improve attentional function for individuals living with moderate to severe TBI (Alvarez et al., 2003; Poon et al., 2020).



KEY POINTS

- Cerebrolysin may be beneficial for improving attentional functioning following brain injury; however, controlled trials are needed to further evaluate its efficacy.

Rivastigmine

Rivastigmine acts as an acetylcholinesterase inhibitor (Liepert, 2016) which prevents the enzyme acetylcholinesterase from breaking down acetylcholine. This increases the concentration of acetylcholine in synapses. Acetylcholine has been most strongly linked with the hippocampus and memory deficits and it is also implicated in attentional processing. Similar to donepezil, rivastigmine has been used to treat dementia in Alzheimer’s disease and to facilitate cognitive recovery in individuals with TBI (Kakehi & Tompkins, 2021).

TABLE 22 | The Effect of Rivastigmine on Attention and Processing Speed Post TBI

Author Year Country Study Design Sample Size	Methods	Outcome
Tenovuo et al. (2009) Finland RCT PEDro=10 N=102	Population: TBI; Mean age=45.5yr; Gender: Males=61, Female=39; Mean time post-injury=8yr; Mean GCS=11. Intervention: Individuals were randomized to receive one of two dosing rivastigmine schedules (placebo then rivastigmine or rivastigmine then placebo). Treatment lasted 8 weeks once a max dose of 12mg per day was reached. Outcome Measures: Computer-based reaction time (CRT), subtraction test, vigilance test (0-5mins, 5-10mins, 10-15mins, correct responses), Symptom Checklist-90 (SC), Diener satisfaction of life scale, Finnish Traumatic Brain Injury Questionnaire (FITBIQ).	<ol style="list-style-type: none"> 1. The percentage of right answers in the subtraction tests were significantly different between groups (p<0.05), and in the third Vigilance test period (OR 0.08; 95% CI 0.001–0.17; p = 0.048) was significantly higher during rivastigmine than placebo treatment (MIXED procedure). 2. Vigilance scores were significantly higher during periods of rivastigmine treatment compared to placebo treatments (p<0.05). 3. There were no other significant differences between groups on any other measures.
Silver et al. (2009)	Population: TBI. <i>Rivastigmine</i> (n=65): Mean Age=36.9yr; Gender: Male=43, Female=22; Time Post	<ol style="list-style-type: none"> 1. The mean final dose of rivastigmine was 7.9 mg/day.

Author Year Country Study Design Sample Size	Methods	Outcome
USA RCT PEDro=9 N=127	<p>Injury=73.5mo. <i>Placebo</i> (n=62): Mean Age=38yr; Gender: Male=42, Female=20; Time Post Injury=100.1mo.</p> <p>Intervention: Participants were randomized to receive rivastigmine injections (1.5 mg 2x/d to a max of 12 mg/d) or placebo injection.</p> <p>Outcome Measures: Trails A and B, Hopkins verbal learning test (HVLTL), Cambridge Neuropsychological Test Automated Batter Rapid Visual Information Processing (CANTAB RVIP A).</p>	<ol style="list-style-type: none"> 40% of patients were responders on CANTAB RVIP A or HVLTL score at week 38. At the end of the study period all (n=98) improved on the CANTAB RVIP A (p<0.001), the HVLTL (P<0.001), and the Trails A and B (p<0.001). Further sub-analysis controlling for order effects demonstrated no significant differences between groups.
Silver et al. (2006) USA RCT PEDro=9 N=123	<p>Population: TBI. <i>Rivastigmine</i> (n=80): Mean Age=37yr; Gender: Male=53, Female=27. <i>Placebo</i> (n=77): Mean Age=37.1yr; Gender: Male=53, Female=24.</p> <p>Intervention: Participants were randomized to receive either rivastigmine (3-6 mg/d) or placebo. At the end of the first 4wk, rivastigmine doses were increased to 3.0 mg, 2x/d. If necessary, doses were decreased to 1.5 mg or 4.5 mg 2x/d.</p> <p>Outcome Measures: Trails A and B, Hopkins verbal learning test (HVLTL), Cambridge Neuropsychological Test Automated Batter Rapid Visual Information Processing (CANTAB RVIP A).</p>	<ol style="list-style-type: none"> Results of the CANTAB RVIP A and HVLTL found no significant differences between the placebo group and the treatment group. Rivastigmine was found to be well tolerated and safe.

Discussion

Three RCTs have concluded that rivastigmine does not improve attention following a TBI (Silver et al., 2006; Silver et al., 2009; Tenovuo et al., 2009). Rivastigmine was safe and well tolerated by individuals with TBI; however, there were no significant differences between rivastigmine and placebo in terms of attention or verbal memory (Silver et al., 2006). In Silver’s (2009) open-label cohort study to their original RCT, participants (n=98) showed significant improvement on the Cambridge Neuropsychological Test Automated Batter Rapid Visual Information Processing (CANTAB RVIP A), the Hopkins verbal learning test (HVLTL) and the trail A and B scales at the end of 38 week study period; however, after further sub-analysis controlling for order effects, no significant differences were found between groups. The study by Tenovuo et al. (2009) found that rivastigmine significantly improved vigilance following doses of 12mg/day for eight weeks. The study by Tenovuo et al. (2009) had higher doses on average and longer duration of rivastigmine administration compared to both Silver et al. studies (Silver et al., 2006; Silver et al., 2009); however, it is unclear whether this resulted in their conflicting results. The route of rivastigmine administration (injection versus oral administration) did not appear to influence its efficacy.

Conclusions

There is level 1a evidence that Rivastigmine compared to placebo is not effective for improving concentration or processing speed in post TBI but may increase vigilance (Silver et al., 2006; Silver et al., 2009; Tenovuo et al., 2009).



KEY POINTS

- Rivastigmine may not be effective in improving concentration or processing speed post TBI.

Amantadine

Amantadine is an agonist of the dopaminergic system and an antagonist of N-methyl-D-aspartate receptors (Liepert, 2016). Amantadine is often used to treat dyskinesia in Parkinson’s disease, as prophylaxis for influenza, and to enhance cognitive recovery in individuals with TBI (Loggini et al., 2020). One study investigated the effect of amantadine on attention in individuals with TBI.

TABLE 23 | Amantadine for the Treatment of Attentional Disorders Following a TBI.

Author Year Country Study Design Sample Size	Methods	Outcome
Hammond et al. (2018) United States RCT PEDro= 9 N=119	<p>Population: TBI; Mean age=38.6yr; Mean time post-injury=6.2yr; Injury severity: GCS<13.</p> <p>Intervention: Individuals were allocated to receive either the placebo or 100mg amantadine 2x/d for 60d. Assessments were completed at baseline, day 28, and day 60.</p> <p>Outcome Measures: Digit-span from Wechsler Memory Scale-III (DS), Trail Making Test (TMT), Controlled Oral Word Association Test (COWAT), Learning/Memory Index (LMI), Attention/Processing Speed Index (APSI), overall composite (GCI).</p>	<ol style="list-style-type: none"> 1. No significant differences were seen on the DS, TMT, COWAT, or the APSI between groups at any time point. 2. The treatment group had significantly lower LMI scores at day 28 compared to the control group (p=0.001), this effect was not present at 60-day follow-up. 3. The treatment group had significantly lower scores on the GCI compared to the control group at day 28 (p=0.002), this effect was not present at day 60 follow-up.

Discussion

Presently, only one study has examined the effects of amantadine on attention and processing speed and found no significant effects following treatment. Any results which were found to be significant on other cognitive measures were not maintained at the 60-day follow-up (Hammond et al., 2018). A recent review on the use of amantadine and memantine to treat individuals with ABI (including stroke), Ma and Zafonte (2020) found that most existing research indicates that amantadine may potentially facilitate cognitive recovery in individuals with ABI; however, the significance of this finding is limited given the

small populations reported by the studies. Research on neurological recovery in populations with ABI is still preliminary and there is limited evidence on risks associated with long-term use of this medication, further studies are needed.

Conclusions

There is level 1b evidence that amantadine is not effective for improving attention compared to placebo following an ABI (Hammond et al., 2018).



KEY POINTS

- Amantadine may not be effective in treating attention deficits following an ABI.

Hyperbaric Oxygen Therapy

Hyperbaric oxygen therapy involves the inhalation of pure oxygen under pressure allowing the lungs to absorb more oxygen per breath. Currently hyperbaric oxygen therapy is used to treat decompression sickness, serious infections, and delayed wound healing as a result of a comorbid illness such as diabetes (The Mayo Clinic, 2019). Hyperbaric oxygen therapy has also been used to treat cognitive difficulties associated with TBI and stroke, such as language and comprehension deficits, as well as memory loss (Gonzalez-Portillo et al., 2019).

TABLE 24 | The Effect of Hyperbaric Oxygen on Attention and Processing Speed Post ABI.

Author Year Country Study Design Sample Size	Methods	Outcome
Hadanny et al. (2018) Israel Case Series N=154	<p>Population: Mean age=42.7yr; Gender: Male=58.4%, Female=43.6%; Mean time post-injury=4.6yr; Injury severity: mild=44.8%, moderate=15.6%, severe=39.6%.</p> <p>Intervention: All individuals received hyperbaric oxygen therapy (HBOT). Sessions consisted of 60-90min of 100% oxygen at 1.5/2 ATA exposure 5d/wk.</p> <p>Outcomes: NeuroTrax software subsets: general, memory, executive functions, attention, information processing speed, visual spatial processing, motor skills.</p>	<ol style="list-style-type: none"> 1. On measures of general cognitive functioning there was a significant increase in scores after HBOT treatment (p<0.0001). 2. Memory scores significantly increased following HBOT treatment (p<0.0001). 3. Executive function scores significantly increased following HBOT treatment (p<0.0001). 4. Attentional scores significantly improved following HBOT treatment (p<0.0001). 5. Information processes speed significantly increased following HBOT treatment (p<0.0001).

Author Year Country Study Design Sample Size	Methods	Outcome
		6. Visual spatial processing significantly improved following HBOT treatment (p=0.005). 7. Motor skills significantly improved following HBOT treatment (p<0.0001).

Discussion

From this single case series, hyperbaric oxygen therapy significantly improved both attention and processing speed following treatment five days a week (Hadanny et al., 2018). General improvements in cognitive functioning and visual processing were also reported (Hadanny et al., 2018). This intervention should be studied more rigorously before firm conclusions are drawn on its efficacy. A recent systematic review found that there is a positive potential for the use of hyperbaric oxygen therapy as a treatment for individuals with TBI, during the acute post-injury period; however, the condition of the individual’s lungs (e.g., the presence of acquired ventilator-associated pneumonia) must be considered before treatment (Daly et al., 2018).

Conclusions

There is level 4 evidence that hyperbaric oxygen therapy may improve both attention and processing speed following an ABI (Hadanny et al., 2018).



KEY POINTS

- Hyperbaric oxygen therapy may improve attention and processing speed following an ABI; however, more research is required.

Dextroamphetamine

Dextroamphetamine is another central nervous stimulant, and like methylphenidate, it is commonly used to treat narcolepsy and attention deficit hyperactivity disorder (Cutler et al., 2022). Dextroamphetamine is a non-catecholamine and sympathomimetic amine that acts as a stimulant.

TABLE 25 | The Effect of Dextroamphetamine on Attention and Engagement Post ABI

Author Year Country Study Design Sample Size	Methods	Outcome
Hart et al. (2018) United States RCT PEDro=10 N=32	<p>Population: TBI; <i>DEX Group (N=17)</i>: Mean age=39.6yr; Gender: Male=11, Female=6; Mean GCS=8.2; Mean time post injury=53.6d. <i>Control Group (N=15)</i>: Mean age=38.7yr; Gender: Male=15, Female=0; Mean GCS=7.5; Mean time post injury=60.2d.</p> <p>Intervention: Participants either received the placebo or 10 mg of dextroamphetamine (DEX). Each treatment was administered once per day for 3wk.</p> <p>Outcomes: Moss Attention Rating Scale, Hopkins Rehabilitation Engagement Rating Scale, Cognitive Failures Questionnaire, Rating Scale of Attentional Behavior, Finger Taping Test, the Symbol Digit Modalities Test, Disability Rating Scale, Agitated Behavior Scale (ABS), Profile of Mood States.</p>	<ol style="list-style-type: none"> 1. There was a significant difference between groups on the ABS ($p=0.04$), with the DEX group demonstrating more agitation over time. 2. No other significant between group differences were found.

Discussion

Based on a single study, it does not appear that dextroamphetamine has any beneficial effects on attention or processing speed following an ABI. However, administration of dextroamphetamine did significantly increase agitation over time.

Conclusions

There is level 1b evidence that dextroamphetamine does not improve attention following an ABI (Hart et al., 2018).

KEY POINTS

- Dextroamphetamine may not be an effective treatment for attentional deficits following an ABI and may increase agitation.

REFERENCES

- Alashram, A. R., Annino, G., Padua, E., Romagnoli, C., & Mercuri, N. B. (2019, August). Cognitive rehabilitation post traumatic brain injury: A systematic review for emerging use of virtual reality technology. *Journal of Clinical Neuroscience*, *66*, 209-219.
<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emexb&AN=2001943605>
https://ocul-uwo.primo.exlibrisgroup.com/openurl/01OCUL_UWO/01OCUL_UWO:UWO_DEFAULT?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emexb&rft.genre=article&rft_id=info:doi/10.1016%2Fj.jocn.2019.04.026&rft_id=info:pmid/31085075&rft.issn=0967-5868&rft.volume=66&rft.issue=&rft.spage=209&rft.pages=209-219&rft.date=2019&rft.jtitle=Journal+of+Clinical+Neuroscience&rft.atitle=Cognitive+rehabilitation+post+traumatic+brain+injury%3A+A+systematic+review+for+emerging+use+of+virtual+reality+technology&rft.aulast=Alashram
- Alvarez, X. A., Sampedro, C., Perez, P., Laredo, M., Couceiro, V., Hernandez, A., Figueroa, J., Varela, M., Arias, D., Corzo, L., Zas, R., Lombardi, V., Fernandez-Novoa, L., Pichel, V., Cacabelos, R., Windisch, M., Alexandre, M., & Moessler, H. (2003, Sep). Positive effects of cerebrolysin on electroencephalogram slowing, cognition and clinical outcome in patients with postacute traumatic brain injury: an exploratory study. *Int Clin Psychopharmacol*, *18*(5), 271-278. <https://doi.org/10.1097/01.yic.0000085765.24936.9a>
- Amos, A. (2002, May). Remediating deficits of switching attention in patients with acquired brain injury. *Brain Inj*, *16*(5), 407-413. <https://doi.org/10.1080/02699050110104435>
- Arciniegas, D. B. (2003, Oct). The cholinergic hypothesis of cognitive impairment caused by traumatic brain injury. *Curr Psychiatry Rep*, *5*(5), 391-399. <https://www.ncbi.nlm.nih.gov/pubmed/13678561>
- Azouvi, P., Couillet, J., Leclercq, M., Martin, Y., Asloun, S., & Rousseaux, M. (2004). Divided attention and mental effort after severe traumatic brain injury. *Neuropsychologia*, *42*(9), 1260-1268.
<https://doi.org/10.1016/j.neuropsychologia.2004.01.001>
- Barnett, M., & Reid, L. (2020). The effectiveness of methylphenidate in improving cognition after brain injury in adults: a systematic review. *Brain Inj*, *34*(1), 1-10. <https://doi.org/10.1080/02699052.2019.1667538>
- Beetz, A., Uvnäs-Moberg, K., Julius, H., & Kotrschal, K. (2012). Psychosocial and psychophysiological effects of human-animal interactions: the possible role of oxytocin. *Frontiers in Psychology*, *3*(234), 1-15.

- Boissonnault, E., Higgins, J., LaGarde, G., Barthelemi, D., Lamarre, C., & Dagher, J. H. (2021). Brain stimulation in attention deficits after traumatic brain injury: a literature review and feasibility study. *Pilot and Feasibility Studies*, 7(1), 1-9. <https://doi.org/http://dx.doi.org/10.1186/s40814-021-00859-3>
- Boman, I. L., Lindstedt, M., Hemmingsson, H., & Bartfai, A. (2004, Oct). Cognitive training in home environment. *Brain Inj*, 18(10), 985-995. <https://doi.org/10.1080/02699050410001672396>
- Bosco, F. M., Parola, A., Angeleri, R., Galetto, V., Zettin, M., & Gabbatore, I. (2018, Nov 2018 2019-01-15). Improvement of communication skills after traumatic brain injury: The efficacy of the cognitive pragmatic treatment program using the communicative activities of daily living. *Archives of Clinical Neuropsychology*, 33(7), 875-888. <https://doi.org/http://dx.doi.org/10.1093/arclin/acy041>
- Cacabelos, R. (2007). Donepezil in Alzheimer's disease: From conventional trials to pharmacogenetics. *Neuropsychiatr Dis Treat*, 3(3), 303-333. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2654795/>
- Campbell, K. A., Kennedy, R. E., Brunner, R. C., Hollis, R. D., Lumsden, R. A., & Novack, T. A. (2018). The effect of donepezil on the cognitive ability early in the course of recovery from traumatic brain injury. *Brain Injury*, 32(8), 972-979.
- Campbell, M. (2000). *Rehabilitation for traumatic brain injury: physical therapy practice in context* (2 ed.). Churchill Livingstone
- Cândido, R. C. F., Menezes de Padua, C. A., Golder, S., & Junqueira, D. R. (2021). Immediate-release methylphenidate for attention deficit hyperactivity disorder (ADHD) in adults (Review). *Cochrane Database of Systematic Reviews*(1).
- Cantor, J., Ashman, T., Dams-O'Connor, K., Dijkers, M. P., Gordon, W., Spielman, L., Tsaousides, T., Allen, H., Nguyen, M., & Oswald, J. (2014, Jan). Evaluation of the short-term executive plus intervention for executive dysfunction after traumatic brain injury: a randomized controlled trial with minimization. *Arch Phys Med Rehabil*, 95(1), 1-9.e3. <https://doi.org/10.1016/j.apmr.2013.08.005>
- Carneiro, A. M. C., Poiani, G. C., Zaninoto, A. L., Lazo Osorio, R., Oliveira, M. D. L., Paiva, W. S., & Zângaro, R. A. (2019). Transcranial Photobiomodulation Therapy in the Cognitive Rehabilitation of Patients with Cranioencephalic Trauma [Article]. *Photobiomodulation, Photomedicine, and Laser Surgery*, 37(10), 657-666. <https://doi.org/10.1089/photob.2019.4683>
- Carswell, A., McColl, M. A., Baptiste, S., Law, M., Polatajko, H., & Pollock, N. (2004). The Canadian Occupational Performance Measure: A research and clinical literature review. *Canadian Journal of Occupational Therapy*, 71(4), 210-222.

- Castel-Lacanal, E., Tarri, M., Loubinoux, I., Gasq, D., de Boissezon, X., Marque, P., & Simonetta-Moreau, M. (2014). Transcranial magnetic stimulation in brain injury. *Annales Francaises d'Anesthesie et de Reanimation*, 33, 83-87.
- Chen, A. J., Novakovic-Agopian, T., Nycum, T. J., Song, S., Turner, G. R., Hills, N. K., Rome, S., Abrams, G. M., & D'Esposito, M. (2011, May). Training of goal-directed attention regulation enhances control over neural processing for individuals with brain injury. *Brain*, 134(Pt 5), 1541-1554.
<https://doi.org/10.1093/brain/awr067>
- Chen, S. H., Thomas, J. D., Glueckauf, R. L., & Bracy, O. L. (1997, Mar). The effectiveness of computer-assisted cognitive rehabilitation for persons with traumatic brain injury. *Brain Inj*, 11(3), 197-209.
<http://www.ingentaconnect.com/content/apl/tbin/1997/00000011/00000003/art00004?token=005219c73f7f0c5a8405847447b496e2f5f73446f554779663e33757e6f4f2858592f3f3b57d640982>
- Cicerone, K. D., Dahlberg, C., Malec, J. F., Langenbahn, D. M., Felicetti, T., Kneipp, S., Ellmo, W., Kalmar, K., Giacino, J. T., Harley, J. P., Laatsch, L., Morse, P. A., & Catanese, J. (2005, Aug). Evidence-based cognitive rehabilitation: updated review of the literature from 1998 through 2002. *Arch Phys Med Rehabil*, 86(8), 1681-1692. <https://doi.org/10.1016/j.apmr.2005.03.024>
- Couillet, J., Soury, S., Lebornec, G., Asloun, S., Joseph, P. A., Mazaux, J. M., & Azouvi, P. (2010, Jun). Rehabilitation of divided attention after severe traumatic brain injury: a randomised trial. *Neuropsychol Rehabil*, 20(3), 321-339. <https://doi.org/10.1080/09602010903467746>
- Cutler, A. J., Suzuki, K., Starling, B., Balakrishnan, K., Komaroff, M., Castelli, M., Meeves, S., & Childress, A. C. (2022). Efficacy and Safety of Dextroamphetamine Transdermal System for the Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents: Results from a Pivotal Phase 2 Study. *Journal of Child and Adolescent Psychopharmacology*, 1-9.
- Dahdah, M. N., Bennett, M., Prajapati, P., Parsons, T. D., Sullivan, E., & Driver, S. (2017). Application of virtual environments in a multi-disciplinary day neurorehabilitation program to improve executive functioning using the Stroop task. *NeuroRehabilitation*, 41(4), 721-734. <https://doi.org/10.3233/nre-172183>
- Dahlin, E., Nyberg, L., Bäckman, L., & Neely, A. S. (2008). Plasticity of executive functioning in young and older adults: Immediate training gains, transfer, and long-term maintenance. *Psychology and Aging*, 23(4), 720-730.
- Daly, S., Thorpe, M., Rockswold, S., Hubbard, M., Bergman, T., Samadani, U., & Rockswold, G. (2018). Hyperbaric Oxygen Therapy in the Treatment of Acute Severe Traumatic Brain Injury: A Systematic Review. *Journal of Neurotrauma*, 35, 623–629.

- Dirette, D. K., Hinojosa, J., & Carnevale, G. J. (1999, Dec). Comparison of remedial and compensatory interventions for adults with acquired brain injuries. *J Head Trauma Rehabil*, 14(6), 595-601. <http://graphics.tx.ovid.com/ovftpdfs/FPDDNCJCLGJLHC00/fs046/ovft/live/gv023/00001199/00001199-199912000-00008.pdf>
- Dockree, P. M., Bellgrove, M. A., O'Keefe, F. M., Moloney, P., Aimola, L., Carton, S., & Robertson, I. H. (2006, Jan). Sustained attention in traumatic brain injury (TBI) and healthy controls: enhanced sensitivity with dual-task load. *Exp Brain Res*, 168(1-2), 218-229. <https://doi.org/10.1007/s00221-005-0079-x>
- Dundon, N. M., Dockree, S. P., Buckley, V., Merriman, N., Carton, M., Clarke, S., Roche, R. A., Lalor, E. C., Robertson, I. H., & Dockree, P. M. (2015, Aug). Impaired auditory selective attention ameliorated by cognitive training with graded exposure to noise in patients with traumatic brain injury. *Neuropsychologia*, 75, 74-87. <https://doi.org/10.1016/j.neuropsychologia.2015.05.012>
- Dvorkin, A. Y., Ramaiya, M., Larson, E. B., Zollman, F. S., Hsu, N., Pacini, S., Shah, A., & Patton, J. L. (2013, Aug 09). A "virtually minimal" visuo-haptic training of attention in severe traumatic brain injury. *J Neuroeng Rehabil*, 10, 92. <https://doi.org/10.1186/1743-0003-10-92>
- Dymowski, A. R., Owens, J. A., Ponsford, J. L., & Willmott, C. (2015). Speed of processing and strategic control of attention after traumatic brain injury. *J Clin Exp Neuropsychol*, 37(10), 1024-1035. <https://doi.org/10.1080/13803395.2015.1074663>
- Dymowski, A. R., Ponsford, J. L., Owens, J. A., Olver, J. H., Ponsford, M., & Willmott, C. (2017). The efficacy and safety of extended-release methylphenidate following traumatic brain injury: a randomised controlled pilot study. *Clinical Rehabilitation*, 31(6), 733-741.
- Fasotti, L., Kovacs, F., Eling, P. A. T. M., & Brouwer, W. H. (2000, 2000/01/01). Time Pressure Management as a Compensatory Strategy Training after Closed Head Injury. *Neuropsychological Rehabilitation*, 10(1), 47-65. <https://doi.org/10.1080/096020100389291>
- Fiani, B., Covarrubias, C., Wong, A., T., D., Reardon, T., Nikolaidis, D., & Sarno, E. (2021). Cerebrolysin for stroke, neurodegeneration, and traumatic brain injury: review of the literature and outcomes. *Neurological Sciences*, 42, 1345-1353.
- Florentino, S. A., Mohammad, H. B., & Ma, H. M. (2022). Acetylcholinesterase inhibitors to enhance recovery from traumatic brain injury: a comprehensive review and case series. *Brain Injury*.
- Foley, J. A., Cantagallo, A., Della Sala, S., & Logie, R. H. (2010). Dual task performance and post traumatic brain injury. *Brain Inj*, 24(6), 851-858. <https://doi.org/10.3109/02699051003789278>

[Record #4181 is using a reference type undefined in this output style.]

- Frenette, A. J., Kanji, S., Rees, L., Williamson, D. R., Perreault, M. M., Turgeon, A. F., Bernard, F., & Fergusson, D. A. (2012). Efficacy and Safety of Dopamine Agonists in Traumatic Brain Injury: A Systematic Review of Randomized Controlled Trials. *Journal of Neurotrauma*, *29*, 1-18.
- Gabbatore, S., K., Angeleri, R., Zettin, M., Bara, B. G., & Bosco, F. M. (2015). Cognitive Pragmatic Treatment: A Rehabilitative Program for Traumatic Brain Injury Individuals. *The Journal of Head Trauma Rehabilitation*, *30*(5), E14-E28.
- Gerber, L. H., Narber, C. G., Vishnoi, N., Johnson, S. L., Chan, L., & Duric, Z. (2014, Aug 08). The feasibility of using haptic devices to engage people with chronic traumatic brain injury in virtual 3D functional tasks. *J Neuroeng Rehabil*, *11*, 117. <https://doi.org/10.1186/1743-0003-11-117>
- Gocheva, V., Hund-Georgiadis, M., & Hediger, K. (2018, Jan 2018 2018-02-15). Effects of animal-assisted therapy on concentration and attention span in patients with acquired brain injury: A randomized controlled trial. *Neuropsychology*, *32*(1), 54-64. <https://doi.org/http://dx.doi.org/10.1037/neu0000398>
- Godfrey, J. (2009, Mar). Safety of therapeutic methylphenidate in adults: a systematic review of the evidence. *J Psychopharmacol*, *23*(2), 194-205. <https://doi.org/10.1177/0269881108089809>
- Gonzalez-Portillo, B., Lippert, T., Nguyen, H., Lee, J. Y., & Borlongan, C. V. (2019). Hyperbaric oxygen therapy: A new look on treating stroke and traumatic brain injury. *Brain Cirulation* *5*(3).
- Gray, J. M., Robertson, I., Pentland, B., & Anderson, S. (1992). Microcomputer-based attentional retraining after brain damage: A randomised group controlled trial. *Neuropsychological Rehabilitation*, *2*(2), 97-115. <http://www.scopus.com/inward/record.url?eid=2-s2.0-0002832362&partnerID=40&md5=12f6d5d007fecb4caf367652889e165c>
- Grealy, M. A., Johnson, D. A., & Rushton, S. K. (1999, Jun). Improving cognitive function after brain injury: the use of exercise and virtual reality. *Arch Phys Med Rehabil*, *80*(6), 661-667. <http://www.sciencedirect.com/science/article/pii/S0003999399901697>
- Gualtieri, C. T., & Evans, R. W. (1988, Oct-Dec). Stimulant treatment for the neurobehavioural sequelae of traumatic brain injury. *Brain Inj*, *2*(4), 273-290.

Hadanny, A., Abbott, S., Suzin, G., Bechor, Y., & Efrati, S. (2018, 28 Sep). Effect of hyperbaric oxygen therapy on chronic neurocognitive deficits of post-traumatic brain injury patients: retrospective analysis. *BMJ Open*, 8(9), e023387.

<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emexa&AN=624242033>

http://vr2pk9sx9w.search.serialssolutions.com/?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emexa&rft.genre=article&rft_id=info:doi/10.1136%2Fbmjopen-2018-023387&rft_id=info:pmid/30269074&rft.issn=2044-6055&rft.volume=8&rft.issue=9&rft.spage=e023387&rft.pages=e023387&rft.date=2018&rft.jtitle=BMJ+open&rft.atitle=Effect+of+hyperbaric+oxygen+therapy+on+chronic+neurocognitive+deficits+of+post-traumatic+brain+injury+patients%3A+retrospective+analysis&rft.aulast=Hadanny

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6169752/pdf/bmjopen-2018-023387.pdf>

Hammond, F. M., Sherer, M., Malec, J. F., Zafonte, R. D., Dikmen, S., Bogner, J., Bell, K. R., Barber, J., & Temkin, N. (2018, 2018 Oct 01

2018-10-18). Amantadine did not positively impact cognition in chronic traumatic brain injury: A multi-site, randomized, controlled trial. *Journal of Neurotrauma*, 35(19), 2298-2305.

<https://doi.org/http://dx.doi.org/10.1089/neu.2018.5767>

Hanh, B., Wolkenberg, F. A., Ross, T. J., Myers, C. S., Heishman, S. J., Stein, D. J., Kurup, P. K., & Stein, E. A. (2008). Divided versus selective attention: evidence for common processing mechanisms. *Brain Res*, 18(1215), 137-146.

Hart, T., Whyte, J., Watanabe, T., & Chervoneva, I. (2018). Effects of dextroamphetamine in subacute traumatic brain injury: A randomized, placebo-controlled pilot study. *Journal of Neuroscience Research*, 96(4), 702–710.

Hasegawa, J., & Hoshiyama, M. (2009, Apr). Attention deficits of patients with chronic-stage traumatic brain injury: a behavioral study involving a dual visuo-spatial task. *J Clin Exp Neuropsychol*, 31(3), 292-301.

<https://doi.org/10.1080/13803390802082054>

Hediger, K., Thommen, S., Wagner, C., Gaab, J., & Hund-Georgiadis, M. (2019, Apr 9). Effects of animal-assisted therapy on social behaviour in patients with acquired brain injury: a randomised controlled trial. *Sci Rep*, 9(1), 5831.

<https://doi.org/10.1038/s41598-019-42280-0>

Hellgren, L., Samuelsson, K., Lundqvist, A., & Borsbo, B. (2015). Computerized Training of Working Memory for Patients with Acquired Brain Injury. *Archives of Physical Medicine and Rehabilitation*, 96(10), e48-e49.

<https://doi.org/10.1016/j.apmr.2015.08.161>

- Huang, C.-H., Huang, C.-C., Sun, C.-K., Lin, G.-H., & Hou, W.-H. (2016). Methylphenidate on Cognitive Improvement in Patients with Traumatic Brain Injury: A Meta-Analysis. *Current Neuropharmacology*, 14.
- Jenkins, P., De Simoni, S., Bourke, N. J., Fleminger, J., Scott, G., Towey, D. J., Svensson, W., Khan, S., Patel, M. C., Greenwood, R., Friedland, D., Hampshire, A., Cole, J. H., & Sharp, D. J. (2019, Aug 1). Stratifying drug treatment of cognitive impairments after traumatic brain injury using neuroimaging. *Brain*, 142(8), 2367-2379. <https://doi.org/10.1093/brain/awz149>
- Jenkins, P. O., De Simoni, S., Bourke, N. J., Fleminger, J., Scott, G., Towey, D. J., Svensson, W., Khan, S., Patel, M. C., Greenwood, R., Friedland, D., Hampshire, A., Cole, J. H., & Sharp, D. J. (2019, Aug 2019 2020-01-30). Stratifying drug treatment of cognitive impairments after traumatic brain injury using neuroimaging. *Brain: A Journal of Neurology*, 142(8), 2367-2379. <https://doi.org/http://dx.doi.org/10.1093/brain/awz149>
- Takehi, S., & Tompkins, D. M. (2021). A Review of Pharmacologic Neurostimulant Use During Rehabilitation and Recovery After Brain Injury. *Annals of Pharmacotherapy*, 55(1254-1266).
- Kang, Y., Gruber, J., & Gray, J. R. (2013). Mindfulness and De-Automatization. *Emotion Review*, 5(2), 192-201.
- Khateb, A., Ammann, J., Annoni, J. M., & Diserens, K. (2005). Cognition-enhancing effects of donepezil in traumatic brain injury. *Eur Neurol*, 54(1), 39-45. <https://doi.org/10.1159/000087718>
- Kim, J., Whyte, J., Patel, S., Europa, E., Wang, J., Coslett, H. B., & Detre, J. A. (2012, Jul). Methylphenidate modulates sustained attention and cortical activation in survivors of traumatic brain injury: a perfusion fMRI study. *Psychopharmacology (Berl)*, 222(1), 47-57. <https://doi.org/10.1007/s00213-011-2622-8>
- Kim, Y. H., Ko, M. H., Na, S. Y., Park, S. H., & Kim, K. W. (2006, Jan). Effects of single-dose methylphenidate on cognitive performance in patients with traumatic brain injury: a double-blind placebo-controlled study. *Clin Rehabil*, 20(1), 24-30. <http://cre.sagepub.com/content/20/1/24.full.pdf>
- Knudsen, E. I. (2007). Fundamental Components of Attention. *Annual Review of Neuroscience* 30(1), 57-78.
- Kullberg-Turtiainen, M., Vuorela, K., Huttula, L., Petri, T., & Sanna, K. (2019). Individualized goal directed dance rehabilitation in chronic state of severe traumatic brain injury: A case study. *Heliyon*, 5(2).
- Lee, S. A., & Kim, M. K. (2018, Dec 4). Effect of Low Frequency Repetitive Transcranial Magnetic Stimulation on Depression and Cognition of Patients with Traumatic Brain Injury: A Randomized Controlled Trial. *Med Sci Monit*, 24, 8789-8794. <https://doi.org/10.12659/msm.911385>

- Lesniak, M. M., Iwanski, S., Szutkowska-Hoser, J., & Seniow, J. (2020, Mar 18). Comprehensive cognitive training improves attention and memory in patients with severe or moderate traumatic brain injury. *Appl Neuropsychol Adult*, 1-10. <https://doi.org/10.1080/23279095.2019.1576691>
- Levin, H., Troyanskaya, M., Petrie, J., Wilde, E. A., Hunter, J. V., Abildskov, T. J., & Scheibel, R. S. (2019, 12 Sep). Methylphenidate Treatment of Cognitive Dysfunction in Adults After Mild to Moderate Traumatic Brain Injury: Rationale, Efficacy, and Neural Mechanisms. *Frontiers in Neurology*, 10 (no pagination), Article 925. <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emexb&AN=629501919>
- https://ocul-uwo.primo.exlibrisgroup.com/openurl/01OCUL_UWO/01OCUL_UWO:UWO_DEFAULT?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emexb&rft.genre=article&rft_id=info:doi/10.3389%2Ffnur.2019.00925&rft_id=info:pmid/&rft.issn=1664-2295&rft.volume=10&rft.issue=&rft.spage=925&rft.pages=&rft.date=2019&rft.jtitle=Frontiers+in+Neurology&rft.atitle=Methylphenidate+Treatment+of+Cognitive+Dysfunction+in+Adults+After+Mild+to+Moderate+Traumatic+Brain+Injury%3A+Rationale%2C+Efficacy%2C+and+Neural+Mechanisms&rft.aulast=Levin
- Levine, B., Robertson, I. H., Clare, L., Carter, G., Hong, J., Wilson, B. A., Duncan, J., & Stuss, D. T. (2000, Mar). Rehabilitation of executive functioning: an experimental-clinical validation of goal management training. *J Int Neuropsychol Soc*, 6(3), 299-312.
- Li, K., Alonso, J., Chadha, N., & Pulido, J. (2015). Does Generalization Occur Following Computer-Based Cognitive Retraining?-An Exploratory Study. *Occup Ther Health Care*, 29(3), 283-296. <https://doi.org/10.3109/07380577.2015.1010246>
- Li, K., Robertson, J., Ramos, J., & Gella, S. (2013, Oct). Computer-based cognitive retraining for adults with chronic acquired brain injury: a pilot study. *Occup Ther Health Care*, 27(4), 333-344. <https://doi.org/10.3109/07380577.2013.844877>
- Liepert, J. (2016). Update on pharmacotherapy for stroke and traumatic brain injury recovery during rehabilitation. *Current Opinion in Neurology*, 29(6).
- Lindelov, J. K., Dall, J. O., Kristensen, C. D., Aagesen, M. H., Olsen, S. A., Snuggerud, T. R., & Sikorska, A. (2016, Oct). Training and transfer effects of N-back training for brain-injured and healthy subjects. *Neuropsychol Rehabil*, 26(5-6), 895-909. <https://doi.org/10.1080/09602011.2016.1141692>
- Loggini, A., Tangonan, R., El Ammar, F., Mansour, A., Goldenberg, F. D., Kramer, C. L., & Lazaridis, C. (2020, July). The role of amantadine in cognitive recovery early after traumatic brain injury: A systematic review.

Clinical Neurology and Neurosurgery, 194 (no pagination), Article 105815.

<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emedx&AN=2005429478>

<https://ocul->

[uwopr.oxfordjournals.org/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emedx&rft.genre=article&rft_id=info:doi/10.1016%2Fj.clineuro.2020.105815&rft_id=info:pmid/&rft.issn=0303-8467&rft.volume=194&rft.issue=&rft.spage=105815&rft.pages=&rft.date=2020&rft.jtitle=Clinical+Neurology+and+Neurosurgery&rft.atitle=The+role+of+amantadine+in+cognitive+recovery+early+after+traumatic+brain+injury%3A+A+systematic+review&rft.aulast=Loggini](https://ocul-uwopr.oxfordjournals.org/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emedx&rft.genre=article&rft_id=info:doi/10.1016%2Fj.clineuro.2020.105815&rft_id=info:pmid/&rft.issn=0303-8467&rft.volume=194&rft.issue=&rft.spage=105815&rft.pages=&rft.date=2020&rft.jtitle=Clinical+Neurology+and+Neurosurgery&rft.atitle=The+role+of+amantadine+in+cognitive+recovery+early+after+traumatic+brain+injury%3A+A+systematic+review&rft.aulast=Loggini)

Ma, H. M., & Zafonte, R. D. (2020, Feb 23). Amantadine and memantine: a comprehensive review for acquired brain injury. *Brain Inj*, 34(3), 299-315. <https://doi.org/10.1080/02699052.2020.1723697>

Magee, W. L., Clark, I., Tamplin, J., & Bradt, J. (2017). Music Interventions for acquired brain injury. *Cochrane Database of Systematic Reviews*(1).

Malec, J., Jones, R. W., Rao, N., & Stubbs, K. (1984). Video game practice effects on sustained attention in patients with craniocerebral trauma. *Cognitive Rehabilitation* 2(4), 18-23.

Manivannan, S., Al-Amri, M., Postans, M., Westacott, L. J., Gray, W., & Zaben, M. (2019, Mar/Apr). The Effectiveness of Virtual Reality Interventions for Improvement of Neurocognitive Performance After Traumatic Brain Injury: A Systematic Review. *J Head Trauma Rehabil*, 34(2), E52-e65. <https://doi.org/10.1097/htr.0000000000000412>

Martinez-Molina, N., Sinponkoski, S., Kuusela, L., Laitinen, S., Holma, M., Ahlfors, M., Jordan-Kilkkki, P., Ala-Kauhala, K., Melkas, S., Pekkola, J., Rodriguez-Fornells, A., Laine, M., Ylinen, A., Rantanen, P., Koskinen, S., Cowley, B. U., & Sarkamo, T. (2021). Resting-State Network Plasticity Induced by Music Therapy after Traumatic Brain Injury. *Neural Plasticity*, 8(6682471), 1-18.

McDowell, S., Whyte, J., & D'Esposito, M. (1998, Jun). Differential effect of a dopaminergic agonist on prefrontal function in traumatic brain injury patients. *Brain*, 121 (Pt 6), 1155-1164.

McHugh, L., & Wood, R. (2013). Stimulus over-selectivity in temporal brain injury: mindfulness as a potential intervention. *Brain Inj*, 27(13-14), 1595-1599. <https://doi.org/10.3109/02699052.2013.834379>

McMillan, T., Robertson, I. H., Brock, D., & Chorlton, L. (2002). Brief mindfulness training for attentional problems after traumatic brain injury: A randomised control treatment trial. *Neuropsychological Rehabilitation*, 12(2), 117-125. <http://www.scopus.com/inward/record.url?eid=2-s2.0-0036246478&partnerID=40&md5=38b009e331e35bd385dc5be05dcca2d0>

- Moraes, T. M., Zaninotto, A. L., Neville, I. S., Hayashi, C. Y., & Paiva, W. S. (2021). Immersive virtual reality in patients with moderate and severe traumatic brain injury: a feasibility study. *Health and Technology, 11*(5), 1035-1044.
- Moseley, A. M., Herbert, R. D., Sherrington, C., & Maher, C. G. (2002). Evidence for physiotherapy practice: A survey of the Physiotherapy Evidence Database (PEDro). *Australian Journal of Physiotherapy, 48*, 43-49.
- Neville, I. S., Zaninotto, A. L., Hayashi, C. Y., Rodrigues, P. A., Galhardoni, R., Ciampi de Andrade, D., Brunoni, A. R., Amorim, R. L. O., Teixeira, M. J., & Paiva, W. S. (2019, Jul 9). Repetitive TMS does not improve cognition in patients with TBI: A randomized double-blind trial. *Neurology, 93*(2), e190-e199. <https://doi.org/10.1212/wnl.00000000000007748>
- Niemann, H., Ruff, R. M., & Baser, C. A. (1990, Dec). Computer-assisted attention retraining in head-injured individuals: a controlled efficacy study of an outpatient program. *J Consult Clin Psychol, 58*(6), 811-817. <https://doi.org/10.1037//0022-006x.58.6.811>
- Novack, T. A., Caldwell, S. G., Duke, L. W., Bergquist, T. F., & Gage, R. J. (1996). Focused versus Unstructured Intervention for Attention Deficits after Traumatic Brain Injury. *The Journal of Head Trauma Rehabilitation, 11*(3), 52-60. http://journals.lww.com/headtraumarehab/Fulltext/1996/06000/Focused_versus_Unstructured_Intervention_for.8.aspx
- Novakovic-Agopian, T., Chen, A. J., Rome, S., Abrams, G., Castelli, H., Rossi, A., McKim, R., Hills, N., & D'Esposito, M. (2011, Sep-Oct). Rehabilitation of executive functioning with training in attention regulation applied to individually defined goals: a pilot study bridging theory, assessment, and treatment. *J Head Trauma Rehabil, 26*(5), 325-338. <https://doi.org/10.1097/HTR.0b013e3181f1ead2>
- O'Neil-Pirozzi, T. M., & Hsu, H. (2016). Feasibility and benefits of computerized cognitive exercise to adults with chronic moderate-to-severe cognitive impairments following an acquired brain injury: A pilot study. *Brain Inj, 30*(13-14), 1617-1625. <https://doi.org/10.1080/02699052.2016.1199906>
- Park, N. W. (1999, 1999/04/01). Evaluation of the Attention Process Training Programme. *Neuropsychological Rehabilitation, 9*(2), 135-154. <https://doi.org/10.1080/713755595>
- Passler, M. A., & Riggs, R. V. (2001). Positive Outcomes in Traumatic Brain Injury–Vegetative State: Patients Treated With Bromocriptine. *Archives of physical medicine and rehabilitation., 82*(3), 311-315.

- Pavlovskaya, M., Hochstein, S., Keren, O., Mordvinov, E., & Groswasser, Z. (2007, May). Methylphenidate effect on hemispheric attentional imbalance in patients with traumatic brain injury: a psychophysical study. *Brain Inj*, 21(5), 489-497. <https://doi.org/10.1080/02699050701311117>
- Plenger, P. M., Dixon, C. E., Castillo, R. M., Frankowski, R. F., Yablon, S. A., & Levin, H. S. (1996, Jun). Subacute methylphenidate treatment for moderate to moderately severe traumatic brain injury: a preliminary double-blind placebo-controlled study. *Arch Phys Med Rehabil*, 77(6), 536-540. <http://www.sciencedirect.com/science/article/pii/S0003999396902919>
- Ponsford, J., Bayley, M., Wiseman-Hakes, C., Togher, L., Velikonja, D., McIntyre, A., Janzen, S., & Tate, R. (2014). INCOG Recommendations for Management of Cognition Following Traumatic Brain Injury, Part II: Attention and Information Processing Speed. *Journal of Head Trauma Rehabilitation*, 29(4), 321-337.
- Poon, W., Matula, C., Vos, P. E., Muresanu, D. F., von Steinbuchel, N., von Wild, K., Homberg, V., Wang, E., Lee, T. M. C., Strilciuc, S., & Vester, J. C. (2020, Feb). Safety and efficacy of Cerebrolysin in acute brain injury and neurorecovery: CAPTAIN I-a randomized, placebo-controlled, double-blind, Asian-Pacific trial. *Neurol Sci*, 41(2), 281-293. <https://doi.org/10.1007/s10072-019-04053-5>
- Roitsch, J., Redman, R., Michalek, A. M. P., Johnson, R. K., & Raymer, A. M. (2019, Jul/Aug). Quality Appraisal of Systematic Reviews for Behavioral Treatments of Attention Disorders in Traumatic Brain Injury. *J Head Trauma Rehabil*, 34(4), E42-e50. <https://doi.org/10.1097/htr.0000000000000444>
- Rosso, M., Buchaim, D. V., Kawano, N., Furlanette, G., Pomini, K. T., & Buchaim, R. L. (2018). Photobiomodulation Therapy (PBMT) in Peripheral Nerve Regeneration: A Systematic Review. *Bioengineering* 5(2), 44.
- Ruff, R., Mahaffey, R., Engel, J., Farrow, C., Cox, D., & Karzmark, P. (1994). Efficacy study of THINKable in the attention and memory retraining of traumatically head-injured patients. *Brain Injury*, 8(1), 3-14. <http://www.scopus.com/inward/record.url?eid=2-s2.0-0028078757&partnerID=40&md5=a6df4d1e06453dcf60ffddcff0fce50>
- Rushby, J. A., De Blasio, F. M., Logan, J. A., Wearne, T., Kornfeld, E., Wilson, E. J., Loo, C., Martin, D., & McDonald, S. (2021, Mar 2). tDCS effects on task-related activation and working memory performance in traumatic brain injury: A within group randomized controlled trial. *Neuropsychol Rehabil*, 31(5), 1-23. <https://doi.org/10.1080/09602011.2020.1733620>
- Sacco, K., Galetto, V., Dimitri, D., Geda, E., Perotti, F., Zettin, M., & Geminiani, G. C. (2016). Concomitant Use of Transcranial Direct Current Stimulation and Computer-Assisted Training for the Rehabilitation of Attention in Traumatic Brain Injured Patients: Behavioral and Neuroimaging Results. *Front Behav Neurosci*, 10, 57. <https://doi.org/10.3389/fnbeh.2016.00057>

- Sackett DL, S. S., Richardson WS, Rosenberg W, Hayes RB. (2000). *Evidence-based medicine: how to practice and teach EBM* (2nd ed. ed.).
- Sarkamo, T., Huttula, L., Hokkanen, L., Koskinen, S., Leppelmeier, J., Saynevirta, K., Molander, K., Forsbom, M.-B., Kullberg-Turtiainen, M., Turtiainen, P., Sarajuuri, J., & Rantanen, P. (2021). DARE to move: feasibility study of a novel dance-based rehabilitation method in severe traumatic brain injury. *Brain Injury*, 35(3), 335-344. <https://doi.org/http://dx.doi.org/10.1080/02699052.2021.1873420>
- Serino, A., Ciaramelli, E., Santantonio, A. D., Malagu, S., Servadei, F., & Ladavas, E. (2007, Jan). A pilot study for rehabilitation of central executive deficits after traumatic brain injury. *Brain Inj*, 21(1), 11-19. <https://doi.org/10.1080/02699050601151811>
- Sigmundsdottir, L., Longley, W. A., & Tate, R. L. (2016). Computerised cognitive training in acquired brain injury: A systematic review of outcomes using the International Classification of Functioning (ICF). *Neuropsychological Rehabilitation*, 26(5-6), 673-741.
- Silver, J. M., Koumaras, B., Chen, M., Mirski, D., Potkin, S. G., Reyes, P., Warden, D., Harvey, P. D., Arciniegas, D., Katz, D. I., & Gunay, I. (2006). Effects of rivastigmine on cognitive function in patients with traumatic brain injury. *Neurology*, 67(5), 748-755. <https://doi.org/10.1212/01.wnl.0000234062.98062.e9>
- Silver, J. M., Koumaras, B., Meng, X., Potkin, S. G., Reyes, P. F., Harvey, P. D., Katz, D. I., Gunay, I., & Arciniegas, D. B. (2009, Feb). Long-term effects of rivastigmine capsules in patients with traumatic brain injury. *Brain Inj*, 23(2), 123-132. <https://doi.org/10.1080/02699050802649696>
- Smart, C. M., Ali, J. I., Viczko, J., & Silveira, K. (2022). Systematic Review of the Efficacy of Mindfulness-Based Interventions as a Form of Neuropsychological Rehabilitation. *Mindfulness*, 13(2), 301–317.
- Sohlberg, M. M., & Mateer, C. A. (2001). Improving attention and managing attentional problems. Adapting rehabilitation techniques to adults with ADD. *Annals of the New York Academy of Sciences*, 931, 359-375.
- Sohlberg, M. M., McLaughlin, K. A., Pavese, A., Heidrich, A., & Posner, M. I. (2000, Oct). Evaluation of attention process training and brain injury education in persons with acquired brain injury. *J Clin Exp Neuropsychol*, 22(5), 656-676. [https://doi.org/10.1076/1380-3395\(200010\)22:5;1-9;ft656](https://doi.org/10.1076/1380-3395(200010)22:5;1-9;ft656)
- Speech, T. J., Rao, S. M., Osmon, D. C., & Sperry, L. T. (1993, Jul-Aug). A double-blind controlled study of methylphenidate treatment in closed head injury. *Brain Inj*, 7(4), 333-338.

- Stablum, F., Umilta, C., Mogentale, C., Carlan, M., & Guerrini, C. (2000). Rehabilitation of executive deficits in closed head injury and anterior communicating artery aneurysm patients. *Psychol Res*, 63(3-4), 265-278.
- Stuss, D. T., Stethem, L. L., Hugenholtz, H., Picton, T., Pivik, J., & Richard, M. T. (1989, Jun). Reaction time after head injury: fatigue, divided and focused attention, and consistency of performance. *J Neurol Neurosurg Psychiatry*, 52(6), 742-748. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1032026/pdf/jnnpsyc00528-0050.pdf>
- Swenson, T. L., Roehmer, C., Tran, R., & Plummer, C. (2021). Donepezil for Aphasia After Severe Traumatic Brain Injury: A Clinical Vignett. *American Journal of Physical Medicine & Rehabilitation*.
- Tenovuo, O., Alin, J., & Helenius, H. (2009). A randomized controlled trial of rivastigmine for chronic sequels of traumatic brain injury-What it showed and taught? *Brain Injury*, 23(6), 548-558. <https://doi.org/10.1080/02699050902926275>
- The Mayo Clinic. (2019). *Hyperbaric Oxygen Therapy*. The Mayo Clinic Retrieved August 8 from <https://www.mayoclinic.org/tests-procedures/hyperbaric-oxygen-therapy/about/pac-20394380>
- Traeger, J., Hoffman, B., Misencik, J., Hoffer, A., & Makii, J. (2020, 01 Apr). Pharmacologic Treatment of Neurobehavioral Sequelae Following Traumatic Brain Injury. *Critical care nursing quarterly*, 43(2), 172-190. <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emexb&AN=631037070>
- https://ocul-uwo.primo.exlibrisgroup.com/openurl/01OCUL_UWO/01OCUL_UWO:UWO_DEFAULT?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emexb&rft.genre=article&rft_id=info:doi/10.1097%2FCNQ.000000000000301&rft_id=info:pmid/32084061&rft.issn=1550-5111&rft.volume=43&rft.issue=2&rft.spage=172&rft.pages=172-190&rft.date=2020&rft.jtitle=Critical+care+nursing+quarterly&rft.atitle=Pharmacologic+Treatment+of+Neurobehavioral+Sequelae+Following+Traumatic+Brain+Injury&rft.aulast=Traeger
- Vester, J. C., Buzoianu, A. D., Florian, S. I., Hömberg, V., Kim, S. H., Lee, T. M. C., C., M., Poon, W. S., Sandesc, D., von Steinbüchel, N., Strilciuc, S., Vos, P. E., von Wild, K., & D., M. (2021). Cerebrolysin after moderate to severe traumatic brain injury: prospective meta-analysis of the CAPTAIN trial series. *Neurological Sciences*, 42, 4531-4541.
- Walz, J. A., Mani, R., Alnawmasi, M. M., & Khuu, S. K. (2021). Visuospatial Attention Allocation as an Indicator of Cognitive Deficit in Traumatic Brain Injury: A Systematic Review and Meta-Analysis. *Frontiers in Human Neuroscience*, 15, 675376–675376.

- Whyte, J., Hart, T., Vaccaro, M., Grieb-Neff, P., Risser, A., Polansky, M., & Coslett, H. B. (2004, Jun). Effects of methylphenidate on attention deficits after traumatic brain injury: a multidimensional, randomized, controlled trial. *Am J Phys Med Rehabil*, 83(6), 401-420.
<http://graphics.tx.ovid.com/ovftpdfs/FPDDNCFBGGDPPC00/fs041/ovft/live/gv012/00002060/00002060-200406000-00001.pdf>
- Whyte, J., Vaccaro, M., Grieb-Neff, P., Hart, T., Polansky, M., & Coslett, H. B. (2008, Feb). The effects of bromocriptine on attention deficits after traumatic brain injury: a placebo-controlled pilot study. *Am J Phys Med Rehabil*, 87(2), 85-99. <https://doi.org/10.1097/PHM.0b013e3181619609>
- Whyte, J. M. D. P., Hart, T. P., Schuster, K. B. S., Fleming, M. P., Polansky, M. S., & Coslett, H. B. M. D. (1997). EFFECTS OF METHYLPHENIDATE ON ATTENTIONAL FUNCTION AFTER TRAUMATIC BRAIN INJURY: A Randomized, Placebo-Controlled Trial1. *American Journal of Physical Medicine & Rehabilitation* November/December, 76(6), 440-450.
<https://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=ovftc&AN=00002060-199711000-00002>
http://vr2pk9sx9w.search.serialssolutions.com/?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:ovftc&rft.genre=article&rft_id=info:doi/&rft_id=info:pmid/&rft.issn=0894-9115&rft.volume=76&rft.issue=6&rft.spage=440&rft.pages=440-450&rft.date=1997&rft.jtitle=American+Journal+of+Physical+Medicine+%26+Rehabilitation&rft.atitle=EFFECTS+OF+METHYLPHENIDATE+ON+ATTENTIONAL+FUNCTION+AFTER+TRAUMATIC+BRAIN+INJURY%3A+A+Randomized%2C+Placebo-Controlled+Trial1.&rft.aulast=Whyte
- Willmott, C., & Ponsford, J. (2009, May). Efficacy of methylphenidate in the rehabilitation of attention following traumatic brain injury: a randomised, crossover, double blind, placebo controlled inpatient trial. *J Neurol Neurosurg Psychiatry*, 80(5), 552-557. <https://doi.org/10.1136/jnnp.2008.159632>
- Willmott, C., Ponsford, J., McAllister, T. W., & Burke, R. (2013). Effect of COMT Val158Met genotype on attention and response to methylphenidate following traumatic brain injury. *Brain Injury*, 27(11), 1281–1286.
- Winters Fisher, A. F. (2019, February). Dance/movement therapy & warrior wellness: A pilot case study. *Arts in Psychotherapy*, 62, 52-60.
<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emexa&AN=2001636363>
https://ocul-uwo.primo.exlibrisgroup.com/openurl/01OCUL_UWO/01OCUL_UWO:UWO_DEFAULT?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:journal&rft_id=info:sid/Ovid:emexa&rft.genre=article&rft_id=info:doi/10.1016%2Fj.aip.2018.11.010&rft_id=info:pmid/&rft.issn=0197-4556&rft.volume=62&rft.issue=&rft.spage=52&rft.pages=52-60&rft.date=2019&rft.jtitle=Arts+in+Psychotherapy&rft.atitle=Dance%2Fmovement+therapy+%26+warrior+wellness%3A+A+pilot+case+study&rft.aulast=Winters+Fisher

- Zhang, L., Plotkin, R. C., Wang, G., Sandel, M. E., & Lee, S. (2004, Jul). Cholinergic augmentation with donepezil enhances recovery in short-term memory and sustained attention after traumatic brain injury. *Arch Phys Med Rehabil*, 85(7), 1050-1055. <http://www.sciencedirect.com/science/article/pii/S0003999304000115>
- Zhang, W. T., & Wang, Y. F. (2017). Efficacy of methylphenidate for the treatment of mental sequelae after traumatic brain injury. *Medicine (Spain)*, 96(25), e6960.
- Zhavoronkova, L. A., Shevtsova, T. P., Pozdneev, A. V., Kuptsova, S. V., & Maksakova, O. A. (2016). Dual-task (postural and cognitive) processing in healthy persons and traumatic brain injury patients. *International Journal of Psychophysiology*, 106, 26-27.
- Zickefoose, S., Hux, K., Brown, J., & Wulf, K. (2013, Jun). Let the games begin: a preliminary study using attention process training-3 and Lumosity brain games to remediate attention deficits following traumatic brain injury. *Brain Inj*, 27(6), 707-716. <https://doi.org/10.3109/02699052.2013.775484>